SEP 14, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.tpxempn

Protocol Citation: Luca Tosti, Christian Conrad 2020. Nuclei isolation from snap frozen human pancreatic tissue using a citric acid buffer. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.tpxempn>

MANUSCRIPT CITATION:
<https://www.biorxiv.org/content/10.1101/733964v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it works for us in different human frozen tissues such as pancreas, lung, brain.

Created: Sep 17, 2018

Last Modified: Sep 14, 2020

Nuclei isolation from snap frozen human pancreatic tissue using a citric acid buffer

In 1 collection

Luca Tosti¹, Christian Conrad²

¹Charité Universitätmedizin Berlin/BIH;

²Digital Health Center at BIH/Charité

The Single Cell Ninjas

Luca Tosti

Charité Universitätmedizin Berlin/BIH

ABSTRACT

A protocol to isolate nuclei from snap-frozen pancreatic tissue and protect RNA from RNase-mediated degradation

MATERIALS

MATERIALS

- ⊗ DTT **Sigma Aldrich Catalog #D0632**
- ⊗ citric acid monohydrate **Sigma-aldrich Catalog #33114**
- ⊗ Sucrose **Sigma Aldrich Catalog #S7903**
- ⊗ Dounce homogenizers **Sigma Catalog #D8938-1SET**
- ⊗ RNase Inhibitor (40 U/μL) **Takara Catalog #634888**
- ⊗ Falcon® 5 mL Round Bottom Polystyrene Test Tube, with Cell Strainer Snap Cap **Corning Catalog #352235**
- ⊗ Countess™ Cell Counting Chamber Slides **Contributed by users Catalog #C10314**
- ⊗ 1M MgCl₂ **Ambion Catalog #AM9530G**
- ⊗ Nuclease-Free Water **Thermo Fisher Scientific Catalog # AM9939**
- ⊗ SUPERase• In™ RNase Inhibitor (20 U/μL) **Thermo Fisher Scientific Catalog #cat# AM2694**
- ⊗ KCl 2M **Contributed by users Catalog #AM9640G**
- ⊗ Tris buffer solution pH 7.5 (1 mol/l) Ultrapure Grade **VWR International**
- ⊗ Hoechst 33258 **Thermo Fisher Scientific Catalog #H3569**
- ⊗ Countess II Automated Cell Counter **Thermo Fisher Scientific Catalog #AMQAX1000**

BEFORE START INSTRUCTIONS

Keywords: single-cell RNA-seq, sNuc-RNA-seq, citric acid buffer, snATAC-seq

- **Cool down table-top centrifuge to 4°C.**
- **Place douncers and pestles on ice/in a fridge to cool them down 0-4°C.**
- **Prepare the following buffers and keep them on ice:**

S25

Reagent	Amount (μL)
Sucrose 1.5M	500.00
Citric Acid 250mM	300.00
Hoechst	3.00
Nuclease-free H2O	2197.00
TOTAL	3000.00

S88

Reagent	Amount (μL)
Sucrose 1.5M	1000.00
Citric Acid 250mM	170.00
Nuclease-free H2O	530.00
TOTAL	1700.00

Resuspension Buffer

Reagent	Amount (μL)
KCl 2M	12.50
MgCl ₂ 1M	3.00
Tris-buffer pH=7.5 1M	50.00
RNaseIn 40U/μL	10.00
DTT (1M)	1.00
SuperaseIn 20U/μL	20.00
Hoechst	1.00
Nuclease-free H2O	902.50
TOTAL	1,000.00

- 1 Add 1 mL of cold S25 buffer to the douncer placed on ice

1 mL S25 buffer

- 2 Transfer the tissue in the douncer, crush it with one stroke of the loose pestle and dislodge the

tissue from the bottom using a P200 tip if necessary

- 3 Incubate the tissue for 5 min on ice

4 °C on ice

00:05:00 incubation

- 4 Crush the tissue with 5 more strokes of the loose pestle and leave to incubate for 5 minutes

4 °C on ice

00:05:00 incubation

- 5 Homogenize the tissue with 3 more strokes of the loose pestle and 5 more strokes with the tight pestle

- 6 Filter through a FACS-tube with cell strainer cap (35 µm)

- 7 Wash filter with 250 µL of S25 buffer

250 µL S25 buffer

- 8 Transfer the nuclei suspension to a 1.5 mL eppendorf tube and centrifuge at 4°C, 500g for 5 minutes

00:05:00 centrifugation

Note

You will probably not see the pellet, but nuclei are on the wall of the tube! Be careful when aspirating, leave a few µL of liquid at the bottom in the next step!

- 9 Resuspend the nuclei in 1 mL of S25 buffer and repeat the centrifugation step as in step 8.

🧴 1 mL S25 buffer

⌚ 00:05:00 centrifugation

- 10** Carefully remove the supernatant, resuspend in 300 μ L S25 buffer and load 10 μ L of nuclei suspension in a Countess™ Cell Counting Chamber Slide. Count the Hoechst-positive nuclei. Use the brightfield to evaluate the amount of debris present in the suspension and the overall integrity of the nuclei. For an example of Countess images, please see the expected results in Step 13.

🧴 300 μ L S25 buffer

Step 10 includes a Step case.

High amount of debris

step case

High amount of debris

If you can see high amount of debris (small fragments in the brightfield view, Hoechst-negative), you can perform a density centrifugation step as described here:

1. Go at the bottom of the nuclei suspension and **gently** add 1 volume (300 μ L) of S88 buffer.
2. Centrifuge at 4°C, 1,000g for 10 minutes.
3. Remove the supernatant - **careful not not aspirate nuclei, better to leave a few microliters at the bottom!**
4. Resuspend in 300 μ L of S25 buffer and check again on the Countess - you should now have a cleaner suspension!
5. Proceed to **Step 10**

🧴 300 μ L S88 buffer

🧴 300 μ L S25 buffer

⌚ 00:10:00 centrifugation

➡ go here 10

- 11** Centrifuge the nuclei at 4°C, 500g for 5 minutes

⌚ 00:05:00 centrifugation

- 12** Resuspend in appropriate volume of Resuspension Buffer. The volume can be determined based on the desired final concentration; 20% of the nuclei might be lost during the previous centrifugation step.

13 Count nuclei and proceed with FACS sorting of the nuclei or directly to the encapsulation in oil droplets.

14

Expected result

A

B

C

Fig. 1: Representative images of bright-field (A), DAPI (B) and merged (C) fields of human pancreas nuclei isolated with the citric acid protocol. Notably, clumps of nuclei are almost completely absent when using this protocol.