VERSION 1

MAR 31, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bd96i99e

Protocol Citation: Sarah Romac 2020. Nucleic acids extraction from single cell using MasterPure Complete DNA and RNA purification (Epicenter). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bd96i99e>

MANUSCRIPT CITATION:

Biard et al. Protist 2015.
<http://dx.doi.org/10.1016/j.protis.2015.05.002>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 26, 2020

Last Modified: Mar 31, 2020

Nucleic acids extraction from single cell using MasterPure Complete DNA and RNA purification (Epicenter) V.1

Sarah Romac¹

¹UMR7144-Roscoff Marine Station

Ecology of Marine Plankton (ECOMAP) team - Roscoff
Tech. support email: phytopk@sb-roscoff.fr

Sarah Romac

UMR7144-Roscoff Marine Station

ABSTRACT

Radiolaria are protists which can't be cultivated. These microorganisms have to be isolated by single-cell for genetic identification and it can be difficult to get clean DNA.

Here we optimized a DNA extraction protocol from protist single-cell. It works very well on single-cell Radiolaria, Foraminifers, but also Ciliates, Dinoflagellates, Diatoms.

MATERIALS

Material and Equipment :

- Glass bent micropipettes.
- Silicone Tubing (Cole-Parmer, Tube silicone platinum LS T17 ref. [FV-96410-17](#))
- StereoMicroscope or binocular microscope.
- Petridishes Diameter 70mm.
- Centrifuge 5417R (Eppendorf)
- Thermomixer Eppendorf

Kit, Reagents and Chemicals :

- **MasterPure Complete DNA and RNA purification** (Epicenter Illumina, ref MC85200)
Enzymes (Proteinase K, RNase, DNase) are stored at -20°C
Other reagents from the kit are stored at room temperature.
- Ethanol absolute, molecular biology quality.
- Propan-2-ol quality molecular biology.
- 0,22µm filtered Seawater or artificial Seawater.
- Ice for enzyme storage during labwork.

Wear labcoat, gloves.

BEFORE START INSTRUCTIONS

Prepare Ethanol 70% from absolute Ethanol : Mix 35 mL of absolute Ethanol with 15 mL nuclease-free water in a Falcon 50mL Store at -20°C.

1. Cell Isolation

- 1 Isolate individually protist cells (at least 50µm in length) using a glass bent micropipette under a binocular microscope.
- 2 Wash each cell in three successive baths of 0.22µm-filtered and sterile seawater.
- 3 Transfer subsequently cells in a 1.5mL sterile microtube.
- 4 Add 30 µL of lysis buffer (Tissue and Cell Lysis Solution from MasterPure™ DNA and RNA Purification Kit, Epicenter) and store at -20°C for barcoding treatment / -80°C for meta-omics treatment.

2. Cell lysis

- 5 Pellet cells by centrifugation (2 min at Vmax), throw the supernatant, let ~25-30 µL of liquid.
- 6 Dilute 1 µL of Proteinase K in 300 µL de lysis solution Tissue et Cellule for each sample. Vortex 10 sec for resuspending cells (facultative).

- 7 Add 300 μ L of mix Proteinase K + lysis solution Tissue et Cellule in each sample. Vortex.
- 8 Incubate 15 min at 65°C , 1000 rpm. Put samples in ice 3-5 min.

3. Total nucleic acids precipitation

- 9 Add 150 μ L MPC reagent to 300 μ L of lysed sample. Vortex.

4+ 500 μ L Propan-2-ol. Mix per inversion.
5-
6
7+ 500 μ L d'EtOH 70%. Don't vortex, mix gently the support. Spin 5min at Vmax ;at 12°C.
8-Discard a maximum of supernatant with precaution, without touching the pellet.
9-Let dry 5-10 min at RT.
10-
10 Spin 10 min at 11 000 g, 12°C. If there is no pellet, add more 25 μ L MPC buffer and spin again 10 min at 11 000 g, 12°C.

11 Transfer the supernatant ion a new clean microtube (1,5 mL), discard the pellet.
(To keep the squeletton, keep the tube with the pellet, add 500 μ L MilliQ Water and store at -20°C).

12 Add 500 μ L of Isopropanol. Mix per inversion. Spin 10min at Vmax, 12°C.

13 Discard the supernatant with precaution, without touching the pellet.

- 14 Add 500 μ L of Ethanol 70%. Don't vortex, mix gently the support. Spin 5min at Vmax ;at 12°C.
- 15 Discard a maximum of supernatant with precaution, without touching the pellet.
- 16 Let dry 5-10 min at room temperature..
- 17 Elute in 25 μ L of TE1x buffer.
Vortex and spin shortly.
Store at - 80°C.