JAN 19, 2020

Optical coherence tomography angiography in age-related macular degeneration

Miklos
Resch¹

¹Semmelweis University of Medical Sciences

Miklos Resch

ABSTRACT

All subjects underwent a comprehensive ophthalmic examination, including best-corrected visual acuity (VA) assessment (ETDRS Charts), slit lamp, and fundus examinations. After pupil dilation, OCT-A was performed, using the AngioVue OCT-A system (RTVue-XR Avanti, OptoVue, Fremont, CA, USA). Scans with the highest resolution were obtained in the central 3×3 mm area, centered on the foveola. The superficial capillary plexus was detected automatically between the internal limiting membrane (ILM) and the inner plexiform layer (IPL); and the deep capillary plexus between the IPL and the outer plexiform layer (OPL). Segmentation errors were manually controlled for each layer to avoid the potential artifacts due to the distortion of anatomy with nAMD.

Foveal thickness was measured in the central 1.0 mm area. Superficial and deep vessel density (VD) was evaluated in the whole image, in the central 1 mm area (fovea), and in the 3 mm ring-shaped parafoveal area. Superficial non-flow area and foveal avascular zone (FAZ) areas were measured, using the built-in AngioAnalytics software (Version ReVue 2018.0.0.18) OptoVue system with an automated segmentation algorithm.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bbiwikfe

Protocol Citation: Miklos Resch 2020. Optical coherence tomography angiography in age-related macular degeneration.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bbiwikfe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jan 19, 2020

Last Modified: Jan 19, 2020

PROTOCOL integer ID:
32054

Keywords: OCT-A