APR 14, 2020

🌐 Oral Glucose Tolerance Test in Mouse

Nancy Smith¹, Mourad Ferdaoussi¹, Haopeng Lin¹,
Patrick E Macdonald¹

¹University of Alberta

CIRTNR2FIC

Nancy Smith
University of Alberta

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.ujjeukn

External link:
<http://www.bcell.org>

Protocol Citation: Nancy Smith, Mourad Ferdaoussi, Haopeng Lin, Patrick E Macdonald 2020. Oral Glucose Tolerance Test in Mouse. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.ujjeukn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 11, 2018

Last Modified: Apr 14, 2020

PROTOCOL integer ID:
16715

ABSTRACT

This protocol details the oral glucose tolerance test in mice. This is performed on C57Bl6, transgenic SENP1 KO and ZMIZ1 KO mouse models. Typically GTT's are done at 12 weeks of age. Following the GTT on chow diet, some mice are put on high fat diet for the next 10 weeks. Subsequent glucose tolerance tests are performed following 2 days, 4 weeks and 8 weeks of high fat diet.

Insulin tolerance test can also be performed. Once the *in vivo* studies are complete, the pancreas or islets are isolated for further *ex vivo* experiments.

MATERIALS

MATERIALS

- ☒ Mouse Diet High Fat (60%) **VWR international Ltd Catalog #F3282**
- ☒ 5LOD Irradiated Pico Lab Rodent Diet **Contributed by users Catalog #6954**
- ☒ Sterile Glucose (50% dextrose) solution 25g/50ml **Contributed by users Catalog #06648(13)**
- ☒ Microvette 100 LH **Sarstedt Catalog #20-1282-100**
- ☒ STELLUX Chemi Rodent Insulin ELISA **Alpco Catalog #80-INSMR-CH01**
- ☒ OneTouch Ultra Blue Blood Glucose Strips **Contributed by users Catalog #L8041261**
- ☒ One Touch Ultra 2 Blood Glucose monitoring system **Contributed by users Catalog #L1540947**
- ☒ 1cc Syringes **BD Biosciences Catalog #B309659**
- ☒ 18G gavage needle- curved **Fine Science Tools Catalog #18061-50**

Keywords: Glucose
Tolerance Test, mouse,
insulin, high fat diet, GTT,
glucose, fasting, oral, gavage

Fasting

- 1 Fasting begins first thing in the morning (around 9am). Fast mice for 4-6 hours before OGTT begins. Transfer mice to clean cage and wire top. Keep water bottles during the fasting period.

Note

- If using High Fat diet (HFD), save food to give back at the end of the OGTT.
- mark tails with a sharpie for easier identification during OGTT.

OGTT

- 2 Weigh mouse to obtain body weight for dose calculations.
Calculate bolus dose of glucose (**1g/kg**).

Note

Weight of mouse (g) x 2 = ul sterile glucose (50% dextrose) solution

*If using HFD for 4 weeks or more, use 0.5g/kg (weight of mouse (g) x 1 = ul sterile glucose (50% dextrose) solution)

☐ GTT- template spreadsheet.xlsx

- 3 Prepare syringes using 1cc syringe and 18G oral gavage needle. Load all syringes with calculated glucose dose and position in front of home cage.
- 4 Prior to the delivery of the sterile glucose (50% dextrose) solution, a time “zero” blood glucose level must be measured. Restrain mouse and clean tail with 70% ethanol/gauze. Extract a small amount of blood from the tail vein onto One Touch Ultra blood glucose strips used with the One Touch Ultra 2 blood glucose monitoring system. Blood can be collected into microvette tubes that can be used to extract plasma which can be assayed for plasma insulin (see step 9 for plasma collection).

Note

Microvette tubes are blood collection tubes with a capillary. Tap tube until all the blood is collected in the tube. Place on ice until plasma collection.

- 5 Administer the correct dose of sterile glucose (50% dextrose) solution using the syringe and gavage needle.
 1. Restraint of the Mouse, avoiding an excessively tight grip of the mouse, as this will induce a panic response. Ensure that the mouse cannot bite you and cannot move its head.
 2. Gravity alone should be used to move the cannula down the esophagus. While gliding the cannula, rotate slightly as the cannula goes past the epiglottis and into the esophagus. The mouse should swallow the cannula, and this can be observed.
 3. Inject the fluid slowly.
 4. The cannula is then gently removed and the animal is returned to its cage.
- 6 Begin timing from the point of successfully delivering the sterile glucose (50% dextrose) solution.
- 7 Test and record the blood glucose level (see step 4) at times 15, 30, 60, 90, and 120 minutes after the initial glucose delivery. Record each value on Glucose Tolerance Test record sheet (template found on Step 2). Record condition of animal throughout and after procedure on the Glucose Tolerance Test record sheet.

Note

Extract a small amount of blood from the tail vein at 15, 30, 60 and 120 minutes as well as testing the blood glucose level.

*skip blood collection at 90 minutes to ensure not to exceed 0.007 ml blood/ g body weight.

- 8 Return food and environmental enrichments to cages.

Plasma collection

- 9 Place tubes into pre-chilled centrifuge. Centrifuge for 2 mins at 2000 rpm.

00:02:00

4 °C

- 10** Remove the callipary-lid and dispose in biohazard bin. Place the base lid and seal the microvette (click position).

- 11** Centrifuge at 4°C for 10 mins at 10 000 rpm.

00:10:00

4 °C

- 12** Using a pipette, transfer the plasma (top transparent layer) to a clean and labelled tube.

Note

Store plasma samples at -20°C until assayed.

- 13** Assay for insulin plasma on a STELLUX Chemi Rodent Insulin ELISA kit by ALPCO. See protocol at:

<https://s3.amazonaws.com/alpco-docs/80/80-INSMR-CH01.pdf>