MAR 16, 2021

Oxford Nanopore sequencing and library construction

Rui Zhang¹

¹BGI-Qingdao, BGI-Shenzhen, Qingdao 266555, China

GigaScience Press

BGI

Hongling Zhou

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.btcwnixe

Protocol Citation: Rui Zhang 2021. Oxford Nanopore sequencing and library construction. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.btcwnixe>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Mar 16, 2021

Last Modified: Mar 16, 2021

PROTOCOL integer ID:
48246

ABSTRACT

The humpback puffer, *Tetraodon palembangensis*, is a species of poisonous freshwater pufferfish mainly distributed in Southeast Asia (Thailand, Laos, Malaysia and Indonesia). The humpback puffer has many interesting biological features, such as inactivity, tetrodotoxin production and body expansion. Here, we reported the first chromosome-level genome assembly of the humpback puffer. The genome size is 362 Mb with ~1.78 Mb contig N50 and ~15.8 Mb scaffold N50. Based on the genome, ~61.5Mb (18.11%) repeat sequences were identified, 19,925 genes were annotated, and 90.01% of these genes could be predicted with function. Finally, a phylogenetic tree of ten teleost fish species was constructed, which suggests that humpback puffer and *T. nigroviridis* shared a common ancestor at 18.1 MYA and diverged from *T. rubripes* at 45.8 MYA. The humpback puffer genome will be a valuable genomic resource to illustrate possible mechanisms of tetrodotoxin synthesis and tolerance.

1 ONT Library preparation and Quality Control

- 1.1** After obtaining the qualified DNA, the large size fraction was selected by automated gel electrophoresis (BluePippin).
- 1.2** Next, the DNA was treated with the end-repair/dA tailing module.
- 1.3** After purification, adapter ligation was performed using ligation sequencing kit ({{kit}}, Oxford Nanopore Technologies).
- 1.4** Finally, DNA library was quantified by Qubit.

Library construction process

2 DNA Sequencing

- 2.1 A certain concentration and volume of DNA library was loaded onto a {{cell_num}} flow cell, which was then transferred to Nanopore {{sequencer}} sequencer for real-time single molecule sequencing.

Nanopore single molecule real-time sequencing