NOV 07, 2018

Oxidative stress

PLOS Genetics

Wen Aw¹

¹z3314717@unsw.edu.au

Wen Aw

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.r69d9h6

External link:

<https://doi.org/10.1371/journal.pgen.1007735>

Protocol Citation: Wen Aw
2018. Oxidative stress.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.r69d9h6>

MANUSCRIPT CITATION:

Aw WC, Towarnicki SG, Melvin RG, Youngson NA, Garvin MR, Hu Y, Nielsen S, Thomas T, Pickford R, Bustamante S, Vila-Sanjurjo A, Smyth GK, Ballard JWO (2018) Genotype to phenotype: Diet-by-mitochondrial DNA haplotype interactions drive metabolic flexibility and organismal fitness. PLoS Genet 14(11): e1007735. doi: [10.1371/journal.pgen.1007735](https://doi.org/10.1371/journal.pgen.1007735)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jul 31, 2018

Last Modified: Nov 07, 2018

PROTOCOL integer ID:
14273

- 1 10 Larvae were dissected on ice and placed in 100uL of ice-cold mitochondrial isolation buffer
- 2 The larvae were homogenized by 80 strokes up and down using sterile Kontes pellet pestles
- 3 The homogenate was pipetted into the syringe (with 2cm² of gauze pad), followed by 400uL of isolation buffer
- 4 Gently press through the gauze into a clean 1.5mL tube
- 5 Filtered homogenate was centrifuged at 1500g for 8 min.
- 6 The resulting brown grey pellet was washed with 200uL of isolation buffer 1 to 2 times to remove the lipid.
- 7 The final mitochondrial pellet was re-suspended in 20uL of mitochondrial isolation buffer

- 8 5uL of mitochondria solution was added into 180uL respiration buffer in 96 well plate and stirred by magnetic stirrer
- 9 Add 5uL of 2.20mM ampliflu red (55uM final conc) + 10uL of 0.02U/uL (20U/mL) horseradish peroxidase (1U mL⁻¹ final conc)
- 10 After the reaction was initiated, substrates were added as follows (allowing a period of stabilization between each step (Pyruvate + L-proline + Malate each 10mM, ADP 5mM, Rotenone 0.5uM and Antimycin A 2.5uM))