VERSION 1

AUG 03, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bjabkian

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PBMC- 03 - TEFF+TREG Isolation from PBMC with “Miltenyi CD4+CD25+ Regulatory T cell Isolation Kit”. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjabkian>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Aug 03, 2020

Last Modified: Aug 03, 2020

🌐 PBMC- 03 - TEFF+TREG Isolation from PBMC with “Miltenyi CD4+CD25+ Regulatory T cell Isolation Kit” V.1

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Farmacologia Medica

ABSTRACT

List of published work using this procedure:

- Kustrimovic, N., Comi, C., Magistrelli, L., Rasini, E., Legnaro, M., Bombelli, R., Aleksic, I., Blandini, F., Minafra, B., Riboldazzi, G., Sturchio, A., Mauri, M., Bono, G., Marino, F., & Cosentino, M. (2018). Parkinson's disease patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17 and Treg in drug-naïve and drug-treated patients. *Journal of neuroinflammation*, 15(1), 205. <https://doi.org/10.1186/s12974-018-1248-8>

MATERIALS

MATERIALS

✂ MACS MultiStand **Miltenyi Biotec Catalog #130-042-303**

✂ EDTA **Sigma Aldrich Catalog #ED2SS**

✂ BSA **Sigma Aldrich Catalog #A2153**

✂ CD4 CD25 Regulatory T Cell Isolation Kit **Miltenyi Biotec Catalog #130-091-301**

✂ LD columns **Miltenyi Biotec Catalog #130-042-901**

✂ MS columns **Miltenyi Biotec Catalog #130-042-201**

✂ MidiMACS Separator **Miltenyi Biotec Catalog #130-042-302**

✂ MiniMACS Separator **Miltenyi Biotec Catalog #130-042-102**

INSTRUMENTATION REQUIRED

Laminar flow hood (**Room PS03**)

BEFORE START INSTRUCTIONS

Make sure that the buffer is **cold (+4°C)** by putting it on **ice** for all the time needed to perform this protocol!

You need to obtaine **TEFF and TREG cells uncontaminated for the subsequent cell culture**, hence make sure you are using **sterile buffers** and **sterile plastic disposables** as well.

Moreover, work under laminar flow hood when you are processing samples (from the beginning to the end of the following procedure).

- 1 Isolate PBMCs according either to the standard protocol from fresh blood or from buffy coat (PBMC- 01a - Isolation of Human PBMC from Buffy Coat, PBMC- 01b - Isolation of Human PBMC from Whole Blood).
- 2 Determine the cell number and viability with the microscope by staining with either Türk or Trypan blue. You can use also Cellometer machine. **(PBMC purity should be $\geq 95\%$ with few contaminant PMNs to prevent clogging of the column).**

For manual cell count use Türk solution for checking purity.

Mix 10 μ l of cell suspension with an equal amount of Türk solution (dilution factor = 2), allow mixture 3 min at room temperature.

Take 10 μ l of the mixture and place it inside a Bürker chamber and view under an optical microscope using 40X magnification.

Count the cells in each square found in the four corners and in the central square (see figure 1 below), including those that lie on the bottom and left-hand perimeters, but not those that lie on the top and right hand perimeters (see figure 2 below).

Total number of cells per ml = mean number of cells x dilution factor x 10⁴ (hemacytometer volume).

Figure 1

The gridded area of the chamber consists of nine 1 mmq squares. These squares are subdivided in three directions; 0.0625 mmq, 0.05 mmq and 0.04 mmq. The central square here in Figure 1 is further subdivided into 0.0025 mmq = 1/25 mmq squares. Count cells in 5 squares as shown.

Figure 2

Concerning those cells that lay on the perimeter of the square, count following this scheme.

For automatic cell count with Cellometer machine use Trypan Blue.

The machine will calculate the n° of cells/ml and the % of viability.

Take 10 µl of cell suspension and add an equal amount of Trypan Blue. Use all the volume to place it in a counting chamber. Place the chamber inside Cellometer and count.

Document	
NAME	SOLUTION- 08 - Türk solution
CREATED BY	Farmacologia Medica
	PREVIEW

Document
NAME
	SOLUTION- 09 - Trypan Blue solution
CREATED BY	
Farmacologia Medica	PREVIEW

Equipment	
Cellometer Auto T4	NAME
Automated cell counter	TYPE
Nexcelom Bioscience	BRAND
EuroClone	SKU

3 OPTIONAL STEP

Sorting of TREG is quite long procedure. Especially in clinical studies with whole blood of enrolled subject, it is possible to stop it after PBMC isolation and counting.

In this case, put cells in a flask with complete medium at a concentration of 1×10^6 cells/mL.

Put the flasks in an incubator (37°C, 5% CO₂), and start sorting procedure the day after.

4 Centrifuge the obtained PBMCs at 1200 x g, Room temperature, 00:05:00 .

Aspirate supernatant completely. (Use 15 mL-conical tube)

Note

Work fast, keep cells cold, and use pre-cooled solutions. This will prevent capping of antibodies on the cell surface and non-specific cell labeling.

Volumes for magnetic labeling indicated in this procedure are for up to 10×10^6 total PBMCs. When working with higher than 10×10^6 cells, scale up all reagent volumes and total volumes accordingly.

For optimal performance it is important to obtain a single-cell suspension before magnetic labeling.

Equipment

Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

- 5 Resuspend the pellet in 100 μ L of cold **SOLUTION- 10** (adjust volumes for 10×10^6 cells).

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

6 Add 10 μL of **CD4+T Cell Biotin-Antibody Cocktail** (adjust volumes for 10×10^6 cells).

7 Mix well and incubate for 00:10:00 at 4 °C

8 Add 20 μL of **Anti-Biotin MicroBeads** (adjust volumes for 10×10^6 cells), mix well and incubate 00:15:00 at 4 °C

9 Add 5 μL of **cold SOLUTION- 10** and centrifuge at 1200 x g, Room temperature, 00:05:00

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

10 Discard the supernatant and resuspend the pellet in 500 μL of **cold SOLUTION- 10**.

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

- 11** Place **LD column** in the magnetic field of suitable MACS Separator (violet, see figures below).

Separator must be attached to the MACS multistand (black) in order to work

- 12 Prepare column by rinsing it with 3 mL of cold **SOLUTION- 10** (trash the effluent).

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

- 13 Apply cell suspension onto the column.

- 14 Collect unlabeled cells that pass through column. Wait until the column reservoir is completely empty.

Wash again **2 times** with 3 mL of cold **SOLUTION- 10** and **1 last time** with 2 mL of the same buffer.

Collect total effluent that is consisting of unlabeled pre-enriched CD4+ cell fraction.

Document
NAME SOLUTION- 10 - TEFF/TREG isolation buffer
CREATED BY Farmacologia Medica	<u>PREVIEW</u>

15 Centrifuge the obtained effluent at 1200 x g, Room temperature, 00:05:00

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

16 Remove supernatant and resuspend cell pellet in 100 µL of cold **SOLUTION- 10** (adjust the volumes for 10×10^6 cells).

Note
<p>Use 15 mL-conical tubes.</p> <p>Volumes for magnetic labeling indicated in this procedure are for an initial starting cell number of up to 10×10^6 total PBMCs. For higher initial cell numbers, scale up all reagent volumes accordingly.</p>

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

17 Add 10 μ L of **CD25 MicroBeads** (adjust volumes for 10×10^6 cells), mix well and incubate 00:15:00 at 4 °C in the dark.

18 Add 5 mL of **cold SOLUTION- 10** and centrifuge at 1200 x g, Room temperature, 00:05:00

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

Equipment

Allegra AVANTI 30

NAME

Centrifuge

TYPE

Beckman Coulter

BRAND

Beckman Italy

SKU

- 19 Remove the supernatant and resuspend the cell pellet in 500 µL of cold **SOLUTION- 10**.

Document

NAME

SOLUTION- 10 - TEFF/TREG isolation buffer

CREATED BY

Farmacologia Medica

PREVIEW

- 20 Place the **MS column** in the magnetic field of a suitable MACS Separator (green, see figures below).

Separator must be attached to the MACS multistand (black) in order to work

- 21 Prepare the column by rinsing with 500 μL of **cold SOLUTION- 10** and trash the effluent.

Document
NAME SOLUTION- 10 - TEFF/TREG isolation buffer
CREATED BY Farmacologia Medica	PREVIEW

22 Apply cell suspension onto the column.

23 Collect the flow-through containing unlabeled negative fraction (T effector cells CD25-).

Wait until the column reservoir is completely empty, wash again **3 times** with **2 µL** of **cold SOLUTION- 10**.

Note

Use 15 mL-conical tube

24 Remove the column from the magnet and place it on a suitable collection tube.

Note

Use 15 mL-conical tube

25 Pipette **1 mL** of **cold SOLUTION- 10** onto the column.

Immediately flush out the magnetically labeled cells by **firmly pushing the plunger into the column**.

The cells that are flushed out are CD25 labeled cells positive fraction (T regulatory cells CD25+).

Document
<div>NAME</div> <div>SOLUTION- 10 - TEFF/TREG isolation buffer</div>
<div>CREATED BY</div> <div>Farmacologia Medica</div>	<div>PREVIEW</div>

26 In order to make sure that collection of cells was complete, **repeat the last step TWO more times.**

27 Centrifuge isolated TEFF and TREG at 1200 x g, Room temperature, 00:05:00

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

28 Resuspend TEFF cells in 1 mL of **SOLUTION- 4** and TREG cells in 0.2 mL of **SOLUTION- 4.**

Count them under microscope or Cellometer machine, according to the appropriate procedure (see step 2 of this protocol).

Document

NAME

SOLUTION- 04 - Wash solution (RPMI/FBS) for PBMC

CREATED BY

Farmacologia Medica

PREVIEW

29 OPTIONAL STEPS

*

If required, it is possible to check the purity of isolated TEFF and TREG.

Proceed as follows:

- Put PBMCs (0.5×10^6 cells), Teff (0.5×10^6 cells) and TREG (at least 0.3×10^6 cells) into 3 different BD Tubes;
- Centrifuge at 1200 x g, Room temperature, 00:05:00
- Remove the supernatant and resuspend the pellet in 50 μ L **PBS 1X** ;
- Add the adequate antibodies such as: **CD4 APC-Cy7** (2.5 μ L , BD cat. n. 557871), **CD25 PE** (10 μ L , Miltenyi cat. n. 120-001-311) and **CD127 AF647** (10 μ L , BD cat. n. 558598) or conjugated to other fluorochromes;
- Incubate for 00:20:00 , in the **dark at RT**;

- Wash with 1 mL of **PBS 1X** and centrifuge 1200 x g, Room temperature, 00:05:00 ;
- Resuspend the pellet in 350 µL **PBS 1X** and leave on ice until FACS acquisition with an appropriate protocol.