VERSION 2

NOV 21, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bpxwmppe

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PBMC- 05 - In Vitro Culture of TEFF+TREG - Cytokine Production by TEFF. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bpxwmppe> Version created by Farmacologia Medica

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 21, 2020

Last Modified: Nov 21, 2020

🌐 PBMC- 05 - In Vitro Culture of TEFF+TREG - Cytokine Production by TEFF V.2

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Farmacologia Medica

ABSTRACT

List of published work using this procedure:

- Kustrimovic, N., Comi, C., Magistrelli, L., Rasini, E., Legnaro, M., Bombelli, R., Aleksic, I., Blandini, F., Minafra, B., Riboldazzi, G., Sturchio, A., Mauri, M., Bono, G., Marino, F., & Cosentino, M. (2018). Parkinson's disease patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17 and Treg in drug-naïve and drug-treated patients. *Journal of neuroinflammation*, 15(1), 205. <https://doi.org/10.1186/s12974-018-1248-8>

MATERIALS

MATERIALS

- ☒ Fetal Bovine Serum (FBS) **EuroClone Catalog #ECS0180L-500 ml**
- ☒ RPMI 1640 **EuroClone Catalog #ECM 0495L- 500 ml**
- ☒ Penicillin/Streptomycin **EuroClone Catalog #ECB3001D - 100 ml**
- ☒ L-Glutamine 100X - 100mL **EuroClone Catalog #ECB3000D**
- ☒ M-Phytohaemagglutinin powder **Sigma Aldrich Catalog #L8902-25 mg**
- ☒ Human Interleukin 2 lyophilized powder research grade **Miltenyi Biotec Catalog #130-097-742**

Instrumentation needed:

Sterile plastic disposables
Laminar Flow Hood
Humidified 37°C, 5% CO₂ incubator

- 1 **Isolate TEFF and TREG** with Miltenyi Kit according to the **protocol PBMC- 03**.
- 2 **Count** both **TEFF** and **TREG** following the appropriate protocol (**CELL COUNT- 02** or **CELL COUNT- 03**).

- 3 Use sterile 96-well round bottom plates.

Note

[Consider that these plates can contain a volume of maximum 250µL]

- 4 Centrifuge TEFF and TREG at 1200 x g, Room temperature, 00:05:00

Equipment

Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

- 5 Resuspend **TEFF** and **TREG** in **SOLUTION- 13** at a **concentration of 1x10⁶/mL**.

Document

NAME

SOLUTION- 13 - Complete culture medium

CREATED BY

Farmacologia Medica

PREVIEW

- 6 According to the experimental design, **activate** a desired number of wells containing **TEFF cells** with **PHA 5µg/ml** (final concentration) and **IL-2 40 ng/mL** (final concentration) by diluting the stock aliquots. Leave also wells of **TEFF unstimulated** (resting control).
- 7 Put **TEFF-CPD labeled cells** and **TREG cells** in the 96-well plate at a **ratio of 1:1** (for example, 0.1×10^6 TEFF + 0.1×10^6 TREG) and activate the cells in the well directly (see step 7 for concentrations): include **1 control co-culture** (not treated with test substance) and **treated co-cultures** (+test substance) according to your experimental design.
- 8 Include also a culture of **resting** and **activated TEFF alone** (for example 0.2×10^6 per well), as control for the subsequent ELISA test.
- 9 Put the plate in a **37 °C** incubator for 48 hours.
- 10 At the end of cell culture, **collect the supernatant** in eppendorf tubes (1.5mL) and **centrifuge them** at **1200 x g, Room temperature, 00:05:00** in order to eliminate any residual cells.

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

11 Collect **supernatants** and **store them at -80°C** until ELISA assay is performed.