VERSION 1

OCT 20, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.6h9hb96

Protocol Citation: Laura Armero 2019. PCR. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.6h9hb96>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 14, 2019

Last Modified: Oct 20, 2019

PROTOCOL integer ID:
 26913

PCR V.1

Laura Armero¹

¹Universidad Complutense de Madrid

AEGIS - Madrid iGEM 2019

Laura Armero

ABSTRACT

Our aim with this protocol is to amplify DNA. This protocol has been optimized has a general amplification

As the quantity of DNA is exponentially increased during the performance of the selection, further modification in the numbers of cycle will be needed to be implemented.

GUIDELINES

We sure to have all the surfaces and materials clean before the start.
 All the procedures must be done in an sterile environment to avoid contamination.

MATERIALS

MATERIALS

Speedy Supreme Green Master Mix **NYZtech Catalog #MB39102**

Agarose (LM-ultrapure grade) **NYZtech Catalog #MB123**

■ Aptamer library (order to IDT)

5` - G TTG CTC GTA TTT AGG GAA TG N<sub>10</sub> ACA CCA GTC TTC ATC CGC TTT<sub>6</sub> - 3`

■ Forward primer (order to IDT):

G TAG GCG AAA₆ - Cy3 - 5`

■ Reserve primer (order to IDT):

5` - BiodTG TTG CTC GTA TTT AGG GAA TG

■ Thermocycler

■ TAE buffer

- 1 Prepare the PCR reaction mixture following the specifications below:

Component	Positive control (V; ul)	Negative control (V; ul)
<i>Template</i>	5	0
<i>Fwd primer</i>	1.25	1.25
<i>Rev primer</i>	1.25	1.25
<i>dH2O</i>	15	20
<i>DMSO</i>	2.5	2.5
<i>Master Mix</i>	25	25

- 2 Perform the amplification in a general thermocycler in the following conditions. Adjust the annealing temperature according to the primers used, and the hotstart to the specifications of your polymerase:

Hot start	95 °	5 min	
Amplification cycles			
Denaturing	95 °	30s	
Annealing	52 °	30s	X 15 cycles
Extension	72 °	30s	
Final extension	72 °	3 min	
Hold	4°		

- 3 Prepare a 3% agarose gel. Load the samples and perform the electrophoresis at 90V for 50 min.
- 4 Remove the gel and observe the bands under UV light.