


Oct 23, 2019

Version 2

PCR V.2

DOI

dx.doi.org/10.17504/protocols.io.8nyhvfw



Claudia Troncone Clemente¹

¹Universidad Complutense de Madrid

AEGIS - Madrid iGEM 2019



Laura Armero

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.8nyhvfw>

Protocol Citation: Claudia Troncone Clemente 2019. PCR. protocols.io <https://dx.doi.org/10.17504/protocols.io.8nyhvfw>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: October 23, 2019

Last Modified: October 23, 2019

Protocol Integer ID: 29112

Keywords: pcr, dna, quantity of dna, amplification, general amplification, protocol

Abstract

Our aim with this protocol is to amplify DNA. This protocol has been optimized has a general amplification

As the quantity of DNA is exponentially increased during the performance of the selection, further modification in the numbers of cycle will be needed to be implemented.

Guidelines

We sure to have all the surfaces and materials clean before the start.

All the procedures must be done in an sterile environment to avoid contamination.

Materials

MATERIALS

⊗ Speedy Supreme Green Master Mix **NZYTEch Catalog #MB39102**

⊗ Agarose (LM-ultrapure grade) **NZYTEch Catalog #MB123**

- Aptamer library (order to IDT)

5` - G TTG CTC GTA TTT AGG GAA TG N₄₀ ACA CCA GTC TTC ATC CGC TTT₆ - 3`

- Forward primer (order to IDT):

G TAG GCG AAA₆ - Cy3 - 5`

- Reserve primer (order to IDT):

5` - BiodTG TTG CTC GTA TTT AGG GAA TG

- Thermocycler
- TAE buffer

Troubleshooting

- 1 Prepare the PCR reaction mixture following the specifications below:

Component	Positive control (V; ul)	Negative control (V; ul)
Template	5	0
Fwd primer	1.25	1.25
Rev primer	1.25	1.25
dH ₂ O	15	20
DMSO	2.5	2.5
Master Mix	25	25

- 2 Perform the amplification in a general thermocycler in the following conditions. Adjust the annealing temperature according to the primers used, and the hotstart to the specifications of your polymerase:

Hot start	95 °	5 min	
Amplification cycles			
Denaturing	95 °	30s	
Annealing	52 °	30s	X 15 cycles
Extension	72 °	30s	
Final extension	72 °	3 min	
Hold	4°		

- 3 Prepare a 3% agarose gel. Load the samples and perform the electrophoresis at 90V for 50 min.



- 4 Remove the gel and observe the bands under UV light.