OCT 15, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.786hrze

Protocol Citation: Janet Standeven 2019. PCR (Toehold) Fragment Protocol for a 25ul reaction.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.786hrze>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol in our group and it is a working PCR protocol.

Created: Oct 15, 2019

Last Modified: Oct 15, 2019

PROTOCOL integer ID:
28670

PCR (Toehold) Fragment Protocol for a 25ul reaction

Janet Standeven¹

¹Lambert iGEM

Lambert Igem

MATERIALS

Q5 reaction buffer- 5ul

dNTPs- 1ul

Primer mix- 2.5ul

- (1.25ul per primer)

Q5 High Fidelity Polymerase- .25ul

Template DNA- *** varies 1000ng<

- Lambert iGEM: 1.5ul

H2O- to 25ul

1 Pipette all reaction agents into pcr tube

- 2 put pcr tube in thermal cycler and set the annealing temperature to 55 degrees celsius
- 3 Annealing temperatures vary depending on DNA
- 4 Set elongation time to 30 seconds
- 5 1 min/2kb for Q5 High Fidelity → may vary