AUG 10, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjibkkan

Protocol Citation: Angel A Justiz-Vaillant 2020. Peptide fragment (579-601 from HIV-gp41) conjugated to keyhole limpet haemocyanin to be used as HIV immunogen.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjibkkan>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 10, 2020

Last Modified: Aug 10, 2020

PROTOCOL integer ID:
 40227

🌐 Peptide fragment (579-601 from HIV-gp41) conjugated to keyhole limpet haemocyanin to be used as HIV immunogen.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

Angel A Justiz-Vaillant
 University of the West Indies St. Augustine

ABSTRACT

Chemical synthesis facilitates the generation of peptides which are exceedingly difficult to express in bacteria, peptide/protein backbone modification, the incorporation of unnatural amino acids, and the production or synthesis of D-proteins. The C-terminal cysteine can be added to the amino acidic sequences of HIV peptides (fragment 579-601 of the HIV-gp41 [1]. These peptide fragments were dimerized by cysteine oxidation with dimethyl-sulfoxide [2] to facilitate their conjugation to keyhole limpet hemocyanin that acts as a carrier protein.

Reference:

1. McPhee DA, Kemp BE, Cumming S, Stapleton D, Gust ID, Doherty RR. Recognition of envelope and tat protein synthetic peptide analogs by HIV positive sera or plasma. *FEBS Lett.* 1988;233(2):393-396. doi:10.1016/0014-5793(88)80468-x.
2. Tam JP, Wu CR, Liu w, Zhang JW (1991) Disulfide bond formation in peptides by dimethyl sulfoxide. Scope and applications. *J Am Chem Soc* 113: 66576662.

GUIDELINES

The Protocol has a high level of reproducibility and has worked for many other HIV peptides.

MATERIALS

MATERIALS

⊗ 10mg KLH (Keyhole Limpet Hemocyanin) (Immunological Grade) **G-Biosciences Catalog #786-088**

⊗ Glutaraldehyde, 50% solution **Bio Basic Inc. Catalog #G0875.SIZE.100ml**

⊗ Peptide 579-601 of HIV-gp41 **Contributed by users**

⊗ Fragment 579-601 of gp41 **Contributed by users**

SAFETY WARNINGS

Glutaraldehyde presents serious side effects including skin irritation, nausea, headache, and shortness of breath.

- 1 These peptide fragment (579-601 from HIV-gp41) is dimerized by cysteine oxidation with dimethyl-sulfoxide. The HIV peptide is dissolved in 5% acetic acid to a final concentration of 5.1 mg/ml.
- 2 The pH of the medium is adjusted to 6 with 1 M (NH₄)₂CO₃.
- 3 Dimethyl-sulfoxide is added to 20% of the final volume, and after four hours at room temperature (RT), the solute is extracted.
- 4 Then, the peptide is dissolved in 3 ml 5% trifluoroacetic acid and precipitated with 35 ml cold ether.
- 5 The precipitate is dialyzed against 1.2 liters of deionized water, pH 7 at 4°C overnight.
- 6 The fragment 579-601 of gp 120 was conjugated by the glutaraldehyde method.
- 7 One mg of keyhole limpet hemocyanin (KLH) is dissolved in 2 ml 0.1 M borate buffer (1.24 g boric acid, 1.90 g sodium tetraborate, pH 10, in 500 mL deionized water).
- 8 In a 20 ml glass tube by gentle stirring 1 µmol of the HIV synthetic peptide and 0.2 mL 0.3%

glutaraldehyde solution (ACS reagent grade, pH 5.5, Sigma-Aldrich) are slowly mixed at RT and left to stand for 2 hrs.

- 9 When a yellow coloration is observed this indicates that the conjugation process is successful.
- 10 To blocking the excess of glutaraldehyde, 0.26 ml of 1 M glycine (Sigma-Aldrich) is added.
- 11 The mix is left for 32 min at RT.
- 12 The HIV-hemocynin conjugate is then dialyzed against 1.3 liters 0.1 M of borate buffer, pH 8.4 through the night at 4°C.
- 13 Then the 0.1 M borate buffer is used to dialyze the preparations for 8 hrs at 4°C.