OCT 01, 2019

🌐 Phage lambda genome concatemerization

Marijn Ceelen¹

¹Wageningen University

iGEM Wageningen 2019

Marijn Ceelen
Wageningen University

ABSTRACT

This protocol is used to create concatemers of phage genomes. This is done by ligating phage lambda genomes using the overhangs at the cos site with T4 ligase.

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.7k3hkyn

Protocol Citation: Marijn Ceelen 2019. Phage lambda genome concatemerization. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.7k3hkyn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Sep 23, 2019

Last Modified: Oct 01, 2019

PROTOCOL integer ID:
28027

1 Add the following ingredients in a microcentrifuge tube.

Ingredient	Amount

Ligation buffer 10X	2 µL
DNA	6 µg
PEG 4000 (40% W/W)	2.5 µL
T4 DNA ligase	1 µL (400 Weiss units)
Water	To 20 µL

2 Incubate at 15 °C overnight