

May 18, 2024

Version 2

Phenocycler-Fusion Staining Protocol For Bone Marrow Tissue V.2

DOI

dx.doi.org/10.17504/protocols.io.5jyl8pzodg2w/v2

Kyung J Ahn¹, Shovik Bandyopadhyay², Anusha Thadi¹, Kai Tan¹

¹Children's Hospital of Philadelphia; ²University of Pennsylvania

CHOP TMC CODEX on F...



Kyung J Ahn

Children's Hospital of Philadelphia

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.5jyl8pzodg2w/v2>

Protocol Citation: Kyung J Ahn, Shovik Bandyopadhyay, Anusha Thadi, Kai Tan 2024. Phenocycler-Fusion Staining Protocol For Bone Marrow Tissue. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.5jyl8pzodg2w/v2> Version created by **Kyung J Ahn**

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited



Protocol status: Working

We use this protocol and it's working

Created: May 18, 2024

Last Modified: August 15, 2025

Protocol Integer ID: 100054

Keywords: HuBMAP, CHOP TMC, CODEX, HTAN, BONE MARROW, staining of ffpe bone marrow tissue, fusion staining protocol for bone marrow tissue, ffpe bone marrow tissue, fusion staining protocol, staining protocol, bone marrow tissue, method for antibody, stepwise protocols for tissue procurement, antigen retrieval, tissue sectioning, antibody, tissue procurement, tissue decalcification, tissue storage, akoya phenocycler, phenocycler, photobleaching step, tissue

Funders Acknowledgements:

National Institutes of Health

Grant ID: U2CCA233285

National Institutes of Health

Grant ID: U54HL156090

Abstract

This protocol describes the method for antibody staining of FFPE bone marrow tissue on slides using CODEX barcoded antibodies. The Akoya Phenocycler-Fusion user manual was modified to include photobleaching steps. Included are the stepwise protocols for tissue procurement, tissue decalcification, tissue sectioning, pre-staining, deparaffinization, antigen retrieval, photobleaching, antibody staining, post-fixation, and tissue storage.

Guidelines

- Use positively charged slides.
- Keep tissue sections from drying during transfer steps

Materials

Sample Kit for PhenoCycler-Fusion[™] (Akoya PN: 7000017)

Troubleshooting



Tissue Procurement

20m

- 1 Femoral head bone marrow tissue obtained from total hip arthroplasty surgery was cut coronally in 4-6 mm thick slices by the Anatomical Pathology Lab at Penn Presbyterian Medical Center.
- 2 A dental biopsy drill was used to harvest 4 mm diameter cylindrical bone marrow tissue samples.

10m

10m

Tissue Fixation

1d

- 3 Bone marrow tissue was fixed in 4% paraformaldehyde at 4°C for 24 hours.

1d

Tissue Decalcification

1w 1d

- 4 Bone marrow tissue was washed 3 times for 30 minutes in PBS at 4°C.
- 5 Bone marrow tissue was decalcified in 0.5M EDTA pH8 for 8 days at 4°C. The EDTA was changed every 2 days.

1h 30m

1w 1d

Tissue Fixation

12h

- 6 After decalcification, samples were washed 5 times for 1 hour in 1 X PBS at 4°C.
- 7 Bone marrow tissue was fixed overnight in 4% paraformaldehyde at 4°C.

5h

12h

FFPE Blocks


- 8 Bone marrow tissue was processed and embedded in paraffin by the Children's Hospital of Philadelphia pathology core.

FFPE Tissue Sectioning

12h

- 9 Prepare a water bath at 40°C.

- 10 Section the bone marrow tissue at a thickness of 5 μm using a microtome. 1m
- 11 Place the sectioned tissue in the water bath for a few seconds until the tissue is completely flat and devoid of folds or wrinkles. 1m
- 12 Place a slide in the water bath and gently lay the tissue on the slide as it is removed from the water bath. Make sure the tissue section is placed in the accessible imaging area (gray in image below) of the slide. 1m



- 13 Place the slide on an angled slide holder and let it dry overnight at room temperature. 12h

Tissue Pre-treatment

1h 5m

- 14 Bake sample slide/s in an incubator at 65°C for 1-3 hours until paraffin thoroughly melts. 1h
- 15 Cool sample slide/s at room temperature for 5 minutes to allow the sample slide/s to cool to room temperature. 5m

Tissue Deparaffinization and Hydration

50m



- 16 Immerse sample slide/s in a coplin jar containing the following reagents for 5 minutes each:

50m

- a.Histochoice
- b.Histochoice
- c.100% Ethanol
- d.100% Ethanol
- e.90% Ethanol
- f.70% Ethanol
- g.50% Ethanol
- h.30% Ethanol
- i.ddH2O
- j.ddH2O

Antigen Retrieval

45m

- 17 In a coplin jar, prepare 50 mls of a 1x citrate buffer. Dilute 10x citrate buffer pH6.0 to 1X citrate buffer in ddH2O.
- 18 Immerse sample slide/s in the coplin jar containing the 1x Citrate buffer and cover the coplin jar with aluminum foil.
- 19 Fill the pressure cooker with 1200 mls of ddH2O. Place the aluminum foil covered coplin jar containing sample slide/s in the pressure cooker.
- 20 Set the pressure cooker to the high-pressure protocol and let the tissue incubate for 20 minutes.
- 21 After incubation in the pressure cooker, release the pressure and carefully remove the coplin jar from the pressure cooker and allow to cool at room temperature for 30 minutes.

30m

15m

Wash Tissue

6m

- 22 Place sample slide/s in a coplin jar containing 50 mls of ddH2O for 2 minutes. All volumes for coplin jar incubations will be 50 mls henceforth in this protocol.
- 23 Place sample slide/s in a second coplin jar filled with ddH2O and incubate for 2 minutes.
- 24 Place sample slide/s in a third coplin jar filled with 1X PBS and incubate for 2 minutes.

2m

2m

2m



Photobleach Tissue

1h 30m

- 25 Submerge the sample slide/s in a 150 mm petri dish containing 50 mls of 4.5% (w/v) H₂O₂ and 20mM NaOH in PBS (bleaching solution).
- 26 Sandwich the petri dish between two broad-spectrum LED light sources for 45 minutes at 4°C. 45m
- 27 After 45 minutes move the sample coverslip(s) into a new petri dish with fresh bleaching solution and photobleach for another 45 minutes at 4°C. 45m

Wash Tissue

10m

- 28 Wash the sample slide/s three times for 2 minutes each in coplin jars containing 1X PBS. 6m
- 29 Immerse the sample slide/s in a coplin jar containing 50 mls of CODEX® Hydration Buffer and incubate for 2 minutes. 2m
- 30 Move sample slide/s to a second coplin jar containing CODEX® Hydration Buffer and incubate for another 2 minutes. 2m

Equilibrate Tissue in Staining Buffer

20m

- 31 Move sample slide/s to a coplin jar containing 50 mls of CODEX® Staining Buffer and incubate for 20-30 minutes. 20m

Preparation of the Antibody Cocktail Solution

- 32 Prepare a stock solution of CODEX® Blocking Buffer to be used for the Antibody Cocktail.

A	B
CODEX® Blocking Buffer	1 Sample
Staining Buffer [μL]	181



	A	B
	N Blocker [μ L]	4.75
	G2 Blocker [μ L]	4.75
	J Blocker [μ L]	4.75
	S Blocker [μ L]	4.75
	Total [μ L]	200

- 33 Add CODEX® Blocking Buffer to a tube designated for Antibody Cocktail Staining Solution. The volume of CODEX® Blocking Buffer to be prepared for each sample slide/s can vary depending on the titer and corresponding volume of each antibody. Adjust volume of CODEX® Blocking Buffer so that the final volume of the Antibody Cocktail Staining Solution is a total of 200 μ L per tissue.
- 34 Add the appropriate volume of each CODEX antibody to the Antibody Cocktail Solution.
- 35 Pipette to mix and briefly spin down the tube.

Tissue Staining

16h

- 36 Remove sample slide/s from the coplin jar containing Staining Buffer and place it on the tray of a humidity chamber.
- 37 Add 195 μ L of the Antibody Cocktail to each of the sample slide(s) and ensure that the liquid covers the entire tissue.
- 38 Incubate overnight in the humidity chamber at 4°C.

16h

Wash Tissue

4m

- 39 After overnight antibody incubation, immerse the sample slide/s in a coplin jar containing 50 mls of CODEX® Staining Buffer for 2 minutes.
- 40 Place sample slide/s in a second coplin jar containing CODEX® Staining Buffer for 2 minutes.

2m

2m



Fix Tissue

10m

- 41 Place sample slide/s in a coplin jar containing 50 mls of 1.6% PFA in CODEX® Storage Buffer and incubate for 10 minutes.

10m

Wash Tissue

1m

- 42 Transfer sample slide/s to a coplin jar containing 50 mls of 1X PBS. Lift and immerse the sample slide/s 2-3 times.
- 43 Transfer sample slide/s to a second coplin jar of 1X PBS and immerse sample slide/s 2-3 times.
- 44 Transfer sample slide/s to a third coplin jar of 1X PBS and immerse sample slide/s 2-3 times.

Ice-cold Methanol Incubation

5m

- 45 Transfer sample slide/s to a coplin jar containing Ice cold methanol and incubate for 5 minutes on ice.

5m

Wash Tissue

1m

- 46 Place a coplin jar of 1x PBS next to the methanol coplin jar containing the sample slide/s. Quickly transfer the sample slide/s from methanol to the coplin jar of 1x PBS and immerse sample slide/s 2-3 times.
- 47 Transfer the sample slide/s to a second coplin jar of 1x PBS and immerse sample slide/s 2-3 times.
- 48 Transfer the sample coverslip to a third coplin jar of 1x PBS and immerse sample slide/s 2-3 times.

Fix Tissue

20m

- 49 Prepare the Final Fixative Solution by diluting 20 µl of the CODEX® Fixative Reagent in 1 ml of 1x PBS (Final Fixative Solution).
- 50 Remove sample slide/s from the coplin jar and place it on the tray of a humidity chamber.



51 Add 200 μ L of Final Fixative Solution to each of the sample slide/s and ensure that the entire tissue is covered in fixative solution.

52 Incubate for 20 mins.

20m

Wash Tissue

1m

53 Remove the sample slide/s from the humidity chamber and place in a coplin jar containing 1x PBS. Lift and immerse the sample slide/s 2-3 times.

54 Move the sample slide/s to a second coplin jar containing 1x PBS and immerse sample slide/s 2-3 times.

55 Move the sample slide/s to a third coplin jar containing 1x PBS and immerse sample slide/s 2-3 times.

Store Tissue

56 Transfer sample slide/s to a coplin jar containing 50 mls of CODEX® Storage Buffer and store at 4°C until imaging.