

Oct 06, 2019

Phylogenetic analysis

 [PLOS One](#)

DOI

dx.doi.org/10.17504/protocols.io.7zihp4e

Norfitriah Mohamed Sohaimi¹, Mohd Hair Bejo¹, Abdul Rahman Omar¹, Aini Ideris¹, Nurulfiza Mat Isa²

¹Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.;

²Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.



Norfitriah Mohamed Sohaimi

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.7zihp4e

External link: <https://doi.org/10.1371/journal.pone.0225863>

Protocol Citation: Norfitriah Mohamed Sohaimi, Mohd Hair Bejo, Abdul Rahman Omar, Aini Ideris, Nurulfiza Mat Isa 2019. Phylogenetic analysis. [protocols.io](https://dx.doi.org/10.17504/protocols.io.7zihp4e) <https://dx.doi.org/10.17504/protocols.io.7zihp4e>

Manuscript citation:

Sohaimi NM, Bejo MH, Omar AR, Ideris A, Isa NM (2019) Molecular characterization of fowl adenovirus isolate of Malaysia attenuated in chicken embryo liver cells and its pathogenicity and immunogenicity in chickens. PLoS ONE 14(12): e0225863. doi: [10.1371/journal.pone.0225863](https://doi.org/10.1371/journal.pone.0225863)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 06, 2019

Last Modified: October 06, 2019

Protocol Integer ID: 28426

Materials

Twenty-eight published FAdVs hexon (Table 1) and fiber (Table 2) gene sequences were obtained from GenBank. Another two reference strains from different genus avian adenovirus was used which are Duck adenovirus (Atadenovirus) and Turkey adenovirus (Siadenovirus). Multiple sequence alignment of nucleotide and deduced amino acid were conducted using BioEdit Version 7.2.5 and Mega Version 5 software. A region of L1 loop in hexon gene with 198 amino acids and entire region of fiber gene was selected for construction of phylogenetic tree. Jones-Taylor-Thorton (JTT) model was used to compute distance matrix using MEGA software and followed by phylogenetic tree construction using Neighbour-joining method with 1000 bootstrap replicates [Juliana et al., 2104].

[Table 1.docx](#)[Table 2.docx](#)

