JUL 12, 2019

Pig ICN recording

Jeffrey Ardell¹

¹University of California, Los Angeles

SPARC

Tech. support email: info@neuinfo.org

Scott John

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.2jugcnw

Protocol Citation: Jeffrey Ardell 2019. Pig ICN recording. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.2jugcnw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 02, 2019

Last Modified: Jul 12, 2019

PROTOCOL integer ID:
22868

ABSTRACT

Yorshire pigs are anesthesized for ICN/ neuromodulation recordings. Monitor hemodynamics

mid-sternotomy

1 Animal: Yorkshire Pig

Surgical Prep:

Sedation Telazol (8mg/kg)

Anesthesia isoflurane (1-2% inhalation), concomitant with intermittent boluses of fentanyl (1–3

µg/kg iv).

Vascular Access: femoral artery (pressure), vein (fluids)

Following completion of the surgery, anesthesia was changed to α-chloralose (50 mg/kg intravenous bolus administration followed by continuous infusion at 10 mg·kg⁻¹·h⁻¹ iv).

- 2 Anesthesia for surgery; 1-2% isoflurane
- 3 Vascular access: femoral artery (pressure) veins (fluid)
- 4 mid-sternotomy: access to heart, great vessels
- 5 Activity generated by neurons in the RAGP (or other ICN) is recorded in situ with a linear micro-electrode array. The multichannel linear micro-electrode array (MicroProbes Inc., Guithersberg, MD, USA), consists of 16 platinum/iridium electrodes (25 µm diameter electrode with an exposed tip of 2 mm; impedance 0.3–0.5 M at 1 kHz), is embedded in the right atrial fat that containing the RAGP such that its tip is placed adjacent to right atrial musculature
- 6 The probe was attached to a flexible lead, allowing the probe to be semi-floating. The density of tissue in the ventral right atrial fat helped to maintain position stability over prolonged periods of time (6–8 h of recording)
- 7 The connecting wires of the multichannel electrode, along with earth and reference wires, were attached to a 16-channel microelectrode amplifier with headstage pre-amplifier (A-M systems, Inc., model 3600; Carlsborg, WA, USA).
- 8 For each channel, filters were set to 300 Hz to 3 kHz, and gain to 5 K. A hook electrode was sewn to the atrial myocardium close to the RAGP to provide a reference right atrial electrogram.
- 9 This atrial electrogram was utilized for determination of atrial rate, duration and characterization

of atrial arrhythmias, including atrial fibrillation, and for identification of the timing of atrial electrical artifacts contained within the neural recording data.

- 10 The 16 microelectrode array signals, along with recorded cardiovascular indices (ECG, right atrium electrogram and haemodynamic data see below), were digitized via a Cambridge Electronics Design (Cambridge, UK; model1401) data acquisition system for off-line analysis. The sampling frequency for neuronal data was 5.26 kHz; it was six times lower (0.877 kHz) for all other signals.
- 11 Neuronal activity was identified as action potentials with signal-to-noise ratios greater than 3:1. The activity generated by individual neuronal somata was identified by the amplitude and configuration (waveform recognition) of these recorded action potentials, using the Spike 2 program. By using these techniques and criteria, action potentials generated by individual somata and/or dendrites, rather than axons of passage, can be recorded for extended periods of time
- 12 anesthesia for recording: alpha-chloralose (50 mg/kg iv bolus, 10 mg/kg/hr iv continuous)
- 13 Hemodynamic recording
Systolic LV pressures were assessed by using a 5-F pigtail, 12-pole conductance-pressure catheter connected to an MPVS Ultra processor (Millar Instruments, Inc, Houston, TX) placed in the left ventricle via carotid artery sheath under ultrasound guidance. Proper catheter position was confirmed by the examination of segmental volume signals. Pressure was continuously monitored and recorded throughout experiments. Increases in LV pressures were noted at stimulation onset, confirming successful stimulation capture. Furthermore, a femoral arterial catheter was also placed and used for systemic arterial pressure monitoring