JAN 06, 2020

Plant material preparation and salt imposition for PlantScreen analysis

Magdalena M Julkowska¹, Mariam Awlia²

¹Boyce Thompson Institute; ²University of Cape Town

Salt Lab KAUST

Mariam Awlia
University of Cape Town

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.4qjgvun

Protocol Citation: Magdalena M Julkowska, Mariam Awlia 2020. Plant material preparation and salt imposition for PlantScreen analysis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.4qjgvun>

MANUSCRIPT CITATION:
[High-Throughput Non-destructive Phenotyping of Traits that Contribute to Salinity Tolerance in Arabidopsis thaliana](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 25, 2019

Last Modified: Jan 06, 2020

ABSTRACT

Reproducible and efficient high-throughput phenotyping approaches, combined with advances in genome sequencing, are facilitating the discovery of genes affecting plant performance. Salinity tolerance is a desirable trait that can be achieved through breeding, where most have aimed at selecting for plants that perform effective ion exclusion from the shoots. To determine overall plant performance under salt stress, it is helpful to investigate several plant traits collectively in one experimental setup. Hence, we developed a quantitative phenotyping protocol using a high-throughput phenotyping system, with RGB and chlorophyll fluorescence (ChlF) imaging, which captures the growth, morphology, color and photosynthetic performance of *Arabidopsis thaliana* plants in response to salt stress.

PROTOCOL integer ID:
25067