

Sep 22, 2019

Plaque assay

DOI

dx.doi.org/10.17504/protocols.io.7i8hkhw



Marijn Ceelen¹

¹iGEM Wageningen

iGEM Wageningen 2019



Marijn Ceelen

Wageningen University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.7i8hkhw

Protocol Citation: Marijn Ceelen 2019. Plaque assay. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7i8hkhw>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: September 22, 2019

Last Modified: September 22, 2019

Protocol Integer ID: 27968

Keywords: Plaque Assay, phage, lambda, T7


Abstract


This is a protocol for the quantification of phage lambda and T7 titers by counting plaques.



Materials

MATERIALS

 Liquid LB medium

 LB agar



- 1 Streak LB plate with E. coli (strain should be susceptible to phage lambda. Strains that include a prophage lambda are likely to be resistant. Strains LE392 and DH10B have been used with this protocol) and incubate overnight at 37 °C
- 2 Pick a colony from this plate and use it to inoculate 10 mL of LB. Incubate this culture at 37 °C until OD reaches 2-3. Overnight culture is recommended.
- 3 Make a range of dilutions. Mix 100 µL of phage dilution with 200 µL of E. coli culture. Then incubate for 10 minutes at room temperature. After 00:10:00 add 3 mL of liquid soft LB agar (LB agar with 0.7% agar) (50 °C).
- 4 Pour the resulting mixture on LB plates (preheat plates at 37 °C). Spread the mixture on the plate by moving it. Make sure to work quickly to avoid clumps of solidified agar.
- 5 Incubate for 00:15:00 at room temperature. Afterwards turn the plates over and incubate overnight at 37 °C .
- 6 Use plates with 30-300 plaques to determine phage concentration. Calculate the Plaque Forming Units (PFU)/mL by the following formula: $PFU/mL = N \times 1/DF \times 1/V$. N is the number of plaques of lysis counted on the plate (expressed as PFU); DF is the dilution factor and V is the volume of phage dilution poured on the plate.