OCT 18, 2020

Plaque PCR

Jiaxin Li¹

¹South China University of Technology

Jiaxin Li

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bni9mch6

Protocol Citation: Jiaxin Li
2020. Plaque PCR.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bni9mch6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Oct 18, 2020

Last Modified: Oct 18, 2020

PROTOCOL integer ID:
43329

- 1 Mark independent plaques on the back of the Petri dish with a black pen. Take 6 points in a plate and try to take points randomly and evenly.

- 2 Take 24 1.5 ml EP tubes.
 - A) Add 200 μ l SM Buffer per tube
 - B) Add 2 μ l chloroform to each tube

Sampling

- 3 Suck out the plaque just spotted with the pipette
- 4 Put into the prepared EP tube.
- 5 Open the metal bath to 95 degrees for lysis of the phage
- 6 Place the EP tube at room temperature for one hour to dissolve the sample sufficiently and mix several times during the period.

PCR system

- 7 KOD FX 20ul system
- 8 Add 18.4 μ l to each tube (1.6 μ l template is needed)
- 9 After an hour, put the EP tube in a metal bath and heat it for 10 minutes.

- 10** Centrifugation for 5 min, 12,000 rpm
- 11** Take 1.6ul supernatant from EP tube and add it to new EP tube.
- 12** Centrifuge the EP tube briefly with a centrifuge.
- 13** Place in PCR instrument.