VERSION 8
AUG 10, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p6kdrw

Protocol Citation: Anneleen Decock 2020. plasma preparation exRNAQC.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.p6kdrw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 16, 2018

Last Modified: Aug 10, 2020

PROTOCOL integer ID:
12204

🌐 plasma preparation exRNAQC V.8

Anneleen
Decock¹

¹Center for Medical Genetics Ghent (CMGG), Cancer Research Institute Ghent (CRIG)

CMGG

Anneleen Decock
Center for Medical Genetics Ghent (CMGG), Cancer Research In...

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

This protocol describes how to prepare different plasma fractions from venous blood draw (10 ml) according to the extracellular RNA quality control study.

GUIDELINES

- In the Center for Medical Genetics Gent (CMGG), plasma preparation is performed in the blood lab. Guidelines on how to handle blood samples and how to work in this lab are described in H5.3-OP2-B1.
- Take pictures of tubes (optional) on a white background, upright position.

Keywords: plasma, platelet-rich plasma, PRP, platelet-poor plasma, PPP, platelet-free plasma, PFP, platelets

Materials

- 1
 - Liquid nitrogen
 - Pipettes and filter tips
 - Centrifuge with swinging bucket rotor (centrifugation speed up to 2500 g (rcf)) and soft brake function, and buckets for blood collection tubes and conical 15 ml tubes
 - Mini centrifuge (for centrifugation of 1.5 ml tubes)
 - Disposable lab coat
 - Nitrile powder-free gloves
 - VERSI-DRY Lab Table Soakers (Nalgene, cat. no. 62080-00)
 - Safe-Lock cup DNA LoBind 1.5 ml PCR clean (Eppendorf, cat.no. 022431021)
 - Conical 15 ml polypropylene tube (Cellstar, cat.no. 188271)
 - Racks for different types of tubes

Centrifugation step 1: 20 min at 400 g (rcf) - preparation of...

- 2
 - Invert tubes 5 times before centrifugation.
 - Spin tubes for 20 min at 400 g (rcf) (without brake), at room temperature. Note time point of centrifugation start.
 - Pipette platelet-rich plasma (PRP) carefully into a new collection tube, leave ± 0.5 cm above the buffy coat (do not disturb the buffy coat).
 - Invert the PRP tube before aliquoting. Aliquot the PRP into LoBind tubes, snap freeze in liquid nitrogen and store at -80°C (note time point of snap freeze), and/or continue to prepare platelet-poor plasma (PPP) and/or platelet pellet.

Centrifugation step 2: 10 min at 800 g (rcf) - preparation of...

- 3
 - Preparation of platelet pellet**
 - Pipette 250 μl PRP into a LoBind tube.
 - Spin the PRP for 10 min at 800 g (rcf) (without brake) to obtain a platelet pellet, at room temperature.
 - Carefully remove the plasma and snap freeze the platelet pellet in liquid nitrogen, and store at -80°C (note time point of snap freeze).
 - Preparation of PPP**
 - Spin the remaining PRP for 10 min at 800 g (rcf) (without brake) to obtain platelet-poor plasma (PPP), at room temperature.
 - Pipette PPP carefully into a new collection tube, leave ± 0.5 cm above pellet (do not disturb pellet).
 - Invert the PPP tube before aliquoting. Aliquot the PPP into LoBind tubes, snap freeze in liquid

nitrogen and store at -80 °C (note time point of snap freeze), and/or continue to prepare platelet-free plasma (PFP).

Centrifugation step 3: 15 min at 2500 g (rcf) - preparation o..

- 4**
 - Spin the PPP for 15 min at 2500 g (rcf) (without brake) to obtain platelet-free plasma (PFP), at room temperature.
 - Pipette PFP carefully into a new collection tube, leave ± 0.5 cm above pellet (do not disturb pellet).
 - Aliquot the PFP into LoBind tubes, snap freeze in liquid nitrogen and store at -80°C (note time point of snap freeze).

Notes

- 5** PRP, PPP and PFP contain approximately 125, 10 and less than 1% of platelet concentration in whole blood ($\sim 300,000$ platelets/ μ l), respectively.