SEP 08, 2017

Plasmid construction and viral infection

PLOS One

Yuejun Wang¹

¹Department of Prosthodontics, Peking University School and Hospital of Stomatology

Yuejun Wang

ABSTRACT

The plasmids were constructed using standard methods; all sequences were verified by appropriate restriction digestion and/or sequencing. Human full-length *IGFBP2* cDNA from ASCs fused to a M2-Flag tag was produced with a standard PCR protocol. This sequence (Flag-*IGFBP2*) was subcloned into the pQCXIN retroviral vector with AgeI and BglII restriction sites. For viral infections, MSCs were plated overnight, then infected with retroviruses in the presence of polybrene (6 µg/mL, Sigma-Aldrich, St. Louis, MO, USA) for 12 h. After 48 h, infected cells were selected with 600 µg/mL G418 for 10 days.

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.iehcbb6](https://doi.org/10.17504/protocols.io.iehcbb6)

External link:
<https://doi.org/10.1371/journal.pone.0184182>

Protocol Citation: Yuejun Wang 2017. Plasmid construction and viral infection. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.iehcbb6>

MANUSCRIPT CITATION:
Wang Y, Liu Y, Fan Z, Liu D, Wang F, Zhou Y (2017) IGFBP2 enhances adipogenic differentiation potentials of mesenchymal stem cells from Wharton's jelly of the umbilical cord *via* JNK and Akt signaling pathways. PLoS ONE 12(8): e0184182. doi: [10.1371/journal.pone.0184182](https://doi.org/10.1371/journal.pone.0184182)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 13, 2017

Last Modified: Nov 09, 2017

PROTOCOL integer ID:
6313

Plasmid construction

- 1 The plasmids were constructed using standard methods; all sequences were verified by appropriate restriction digestion and/or sequencing.
- 2 Human full-length *IGFBP2* cDNA from ASCs fused to a M2-Flag tag was produced with a standard PCR protocol.
- 3 This sequence (Flag-*IGFBP2*) was subcloned into the pQCXIN retroviral vector with AgeI and BglII restriction sites.

viral infections

- 4 MSCs were plated overnight, then infected with retroviruses in the presence of polybrene (6 µg/mL, Sigma-Aldrich, St. Louis, MO, USA) for 12 h.
- 5 After 48 h, infected cells were selected with 600 µg/mL G418 for 10 days.