

Sep 08, 2017

Plasmid construction and viral infection

 [PLOS One](#)

DOI

dx.doi.org/10.17504/protocols.io.iehcbb6

Yuejun Wang¹

¹Department of Prosthodontics, Peking University School and Hospital of Stomatology



Yuejun Wang

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.iehcbb6

External link: <https://doi.org/10.1371/journal.pone.0184182>

Protocol Citation: Yuejun Wang 2017. Plasmid construction and viral infection. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.iehcbb6>

Manuscript citation:

Wang Y, Liu Y, Fan Z, Liu D, Wang F, Zhou Y (2017) IGFBP2 enhances adipogenic differentiation potentials of mesenchymal stem cells from Wharton's jelly of the umbilical cord *via* JNK and Akt signaling pathways. PLoS ONE 12(8): e0184182. doi: [10.1371/journal.pone.0184182](https://doi.org/10.1371/journal.pone.0184182)

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: June 12, 2017

Last Modified: November 09, 2017

Protocol Integer ID: 6313



Abstract

The plasmids were constructed using standard methods; all sequences were verified by appropriate restriction digestion and/or sequencing. Human full-length *IGFBP2* cDNA from ASCs fused to a M2-Flag tag was produced with a standard PCR protocol. This sequence (Flag-*IGFBP2*) was subcloned into the pQCXIN retroviral vector with *AgeI* and *BglII* restriction sites. For viral infections, MSCs were plated overnight, then infected with retroviruses in the presence of polybrene (6 µg/mL, Sigma-Aldrich, St. Louis, MO, USA) for 12 h. After 48 h, infected cells were selected with 600 µg/mL G418 for 10 days.



Plasmid construction

- 1 The plasmids were constructed using standard methods; all sequences were verified by appropriate restriction digestion and/or sequencing.
- 2 Human full-length *IGFBP2* cDNA from ASCs fused to a M2-Flag tag was produced with a standard PCR protocol.
- 3 This sequence (Flag-*IGFBP2*) was subcloned into the pQCXIN retroviral vector with AgeI and BglII restriction sites.

viral infections

- 4 MSCs were plated overnight, then infected with retroviruses in the presence of polybrene (6 µg/mL, Sigma-Aldrich, St. Louis, MO, USA) for 12 h.
- 5 After 48 h, infected cells were selected with 600 µg/mL G418 for 10 days.