DEC 15, 2020

Plasmodium berghei ookinete culture

In 1 collection

Benito Recio Totoro^{1,2}

¹Centro de Investigaciones Sobre Enfermedades Infecciosas, Instituto Nacional de Salud Pública;

²Instituto de Biotecnología, Universidad Nacional Autónoma de México

Benito Recio Totoro

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bmm8k49w

Protocol Citation: Benito Recio Totoro 2020. *Plasmodium berghei* ookinete culture. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bmm8k49w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Sep 22, 2020

Last Modified: Dec 15, 2020

PROTOCOL integer ID:
42400

Keywords: *Plasmodium berghei*, Ookinete culture

DISCLAIMER

This protocol involves working with mice which, by the nature of the procedure, will end being sacrificed.

ABSTRACT

Plasmodium berghei is one of the main non-human malaria models used in research. It readily infects laboratory strains of mice and rats as well as mosquitoes. Because of this, it is safe to manipulate and it does not require human fluids/tissues. This document provides the Operational Procedure for producing and culturing ookinetes of *P. berghei in vitro*. The method has been optimized for the *P. berghei* ANKA train clone 2.34 and the derived GFP-expressing clone, but should work for other strains.

This method is adapted and modified from:

CITATION

Rodríguez MC, Margos G, Compton H, Ku M, Lanz H, Rodríguez MH, Sinden RE (2002). *Plasmodium berghei*: routine production of pure gametocytes, extracellular gametes, zygotes, and ookinetes. *Experimental parasitology*.

CITATION

Carter, V., Cable, H. C., Underhill, B. A., Williams, J., & Hurd, H. (2003). Isolation of *Plasmodium berghei* ookinetes in culture using Nycodenz density gradient columns and magnetic isolation.. *Malaria Journal*.

LINK

<https://dx.doi.org/10.1186%2F1475-2875-2-35>

CITATION

Nacer A, Underhill A, Hurd H (2008). The microneme proteins CTRP and SOAP are not essential for *Plasmodium berghei* ookinete to oocyst transformation in vitro in a cell free system.. *Malaria journal*.

LINK

<https://doi.org/10.1186/1475-2875-7-82>

GUIDELINES

Although *Plasmodium berghei* ookinetes have been successfully cultured for more than 30 years, several improvements have been made to the basic protocol. One of them is the use of CF11 cellulose powder to remove white blood cells from the

culture. This is advisable since it has been shown that the white blood cells can induce programmed cell death in the ookinetes. Furthermore, it is also a recommended step if DNA or RNA has to be extracted from the culture because eliminates nucleic acid contamination from the white blood cells. However, we have found the use of CF11 columns troublesome since, in our hands, it causes the culture yield to decrease (Fig. 1) probably because of a reduction in gametocyte viability.

Figure 1. Decrease in yield after passing the infected blood through CF11 cellulose powder columns to eliminate white blood cells. Error bars represent the standard deviation of the mean, n = number of cultures, p = two-tailed t-test.

Working schedule to obtain ookinetes every week.

	Sun.	Mon.	Tues.	Wed.	Thur s.	Fri.	Sat.	
9:00-10:00			Ookinetes ready to use					
10:00-12:00			Phenylhydr azine			Parasite pass for ook. culture and for next week's pass.		
12:00-14:00								

14:00-16:00		Ookinete culture							
-------------	--	------------------	--	--	--	--	--	--	--

BEFORE START INSTRUCTIONS

Requirements:

- 1.- *Plasmodium berghei* cryopreserved stock.
- 2.- Laboratory mice (we use 6 to 8 weeks old male BALB/c mice).
- 3.- CO₂ euthanasia chamber.
- 4.- Brightfield microscope.
- 5.- Microscope slides and coverslips.
- 6.- Newbauer chamber (haemocytometer).
- 7.- 0.6 and 1.5 ml microcentrifuge tubes.
- 8.- Incubator at 20°C (Temp. range 18 to 21°C).
- 9.- 1 ml syringes.
- 10.- Cotton balls.
- 11.- Bunsen burner.
- 12.- T25 culture flasks.
- 13.- Common cell culture labware.

Reagents:

- 1.- Saline solution (0.9%).
- 2.- Methanol.
- 3.- Ethanol (70%).
- 4.- Giemsa stain (30% in water).
- 5.- Immersion oil.
- 6.- Phosphate-buffered-saline (PBS).
- 7.- Phenylhydrazine (6 mg/ml of saline).
- 8.- Heparin (250 IU/ml of saline).
- 9.- Ookinete culture medium: RPMI 1640 medium at pH 8.3 supplemented with 23.81 mM sodium bicarbonate, 0.37 mM hypoxanthine, 25 mM HEPES, 5000 U/ml penicillin, 5 mg/ml streptomycin, 10 mg/ml neomycin, and 20% heat-inactivated fetal bovine serum.

Mice infections for passages.

- 1 Mice are inoculated via intraperitoneally (IP) with no more than 200 µL either from a cryopreserved stock or by passage from other mice.
From cryopreserved stock: allow the blood-cryopreservation solution to thaw slowly and inoculate 200 µL IP.
Passage from another mouse: The number of parasites per ml of blood has to be determined to always inoculate a constant amount.

- 2 To determine the number of parasites per ml of blood, collect a drop of tail blood from the infected mouse and make a thin smear. Let it dry and fix with methanol.
- 3 Prepare 1 mL of 30% Giemsa per slide and stain for 00:10:00 .
- 4 Rinse slide with distilled water and dry by gently applying pressure with an absorbent paper towel. Do not rub the paper towel on the slide.
- 5 Determine the parasitemia by counting at least 500 red blood cells (RBC) at 1000x magnifications and determine the percentage of infected RBC.

$$\text{parasitemia} = (\text{iRBC} / \text{RBC}) (100)$$

where *iRBC* = infected RBC counted, *RBC* = total number of RBC counted.

Note

Infected RBC with more than one parasite count as one.

- 6 Collect an additional 1 μL of tail blood and dilute it to 1 mL with PBS with 1 μL of heparin.
- 7 Mix well and determine the number of RBC per ml of blood with the Neubauer chamber.
 - 7.1 Fill the two slots of the Neubauer chamber with $\sim$ 10 μL of blood solution (which is diluted 1/1000). Do not overflow the chambers.
 - 7.2 View under the microscope first with the 10x objective to locate the grid, and then with a 40x

objective to zoom into the centre grid (the one with the smallest squares). The centre square is a 5 by 5 grid, each of these smaller squares are a 4 by 4 grid. Count the number of RBC in 5 squares of the 5 by 5 grid (the 4 squares in the corners and the centre one). Cells touching the top and right margins are counted, while cells touching the bottom and left margins are not counted.

7.3
$$\text{RBC/ml} = (\text{xRBC}/20 \times 10^6) (1 \times 10^{12}) (1000)$$

8 Determine the number of infected RBC per ml of blood using the parasitemia obtained previously.
$$\text{iRBC/ml} = (\text{parasitemia}/100)(\text{RBC/ml})$$

9 Euthanize the mouse in the CO₂ chamber and collect the blood via cardiac puncture with a 1 ml syringe preloaded with 10 µL of heparin. Before puncture, clean the puncture site with a cotton ball soaked in 70% ethanol.

10 For maintenance passages, dilute the blood with PBS to obtain 4 x 10⁵ infected RBC per 200 µL and inoculate IP the mice.

Note

Make no more than 8 passages as gametocytes lose the ability to mature into gametes. Also, keep infected mice for no more than a week as they will suffer unnecessarily. With this inoculum, the parasitemia should reach between 15 to 25% in one week.

Mice infections for ookinete culture

11 For ookinete culture, inoculate mice with 200 µL of phenylhydrazine IP three days before the inoculation of the parasites to induce reticulocytosis.

12 Determine the parasitemia and infected RBC concentration of the donor mouse as explained in steps 2 to 8.

13 Collect the infected blood as explained in step 9.

14 Dilute the infected blood with PBS to obtain 4×10^7 infected RBC per 200 μL and inoculate the donor mouse IP.

Note

With this inoculum, the parasitemia must reach between 15 to 25% in three days.

Ookinete culture

15 Three days after the inoculum of 4×10^7 infected RBC verify the parasitemia as explained in steps 2 to 5, and verify the viability of the gametocytes by performing an exflagellation test.

15.1 For the exflagellation test, collect an additional 2 μL of tail blood and place in a 0.6 ml microcentrifuge tube with 7 μL of ookinete medium and 1 μL of heparin. Mix very gently. Take the resulting 10 μL and place them on a microscope slide, cover with a coverslip and incubate for 00:15:00 at 20 $^{\circ}\text{C}$.

15.2 Determine the number of exflagellation centres in 10 random fields of view.

Note

We only use mice with 15 to 25% parasitemia and >7 exflagellation centres per field of view.

We find it not useful to determine the gametocytaemia because the correlation of parasitemia vs gametocytaemia, and the correlation between gametocytaemia vs exflagellation is low (Fig 2).

Figure 2. Correlation between parasitemia and gametocytaemia (A), gametocytaemia and exflagellation (B), and exflagellation and number of ookinetes obtained (C).

16

Prepare a T25 culture flask with 4 ml of ookinete medium. Sacrifice the mice and collect the blood as explained in step 9.

Note

About 1 ml of blood is obtained.

It is critical to collect the blood in a gentle manner but fast. Do not pull the syringe plunger too much as to generate vacuum. Mix the blood with the heparin very well. Try to succeed in the cardiac puncture in the first attempt.

Rough manipulation of the blood at this step decreases considerably the ookinete culture yield.

17

Add the blood to the culture flask. Again treat the blood gently, do not push the syringe plunger too hard.

Mix slightly and incubate at 20 °C for 18:00:00.

18

After incubation, determine the number of ookinetes in the culture by taking a sample of

50 μL . Dilute in 950 μL of PBS and count in the Neubauer chamber.

*

18.1

Add ~ 10 μL of the ookinete culture dilution on each chamber and count the number of ookinetes in the four big squares in the corners. For reference, these big squares have a 4 by 4 grid.

$\text{Ookinetes/ml} = (\text{xOokinetes}/4\text{squares})(20)(10000)$

This is the number of ookinetes over the number of squares counted, times the dilution factor, times a conversion factor.

Note

Usually, between 25 and 35 million ookinetes are obtained in total per culture (from one mouse).