NOV 21, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bpxxmppn

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PMN- 01a - Isolation of Human PMN from Buffy Coat. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bpxxmppn>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 21, 2020

Last Modified: Nov 21, 2020

PROTOCOL integer ID:
 44759

PMN- 01a - Isolation of Human PMN from Buffy Coat

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese)

Farmacologia Medica

ABSTRACT

Separation of Human Neutrophils (PMN) from Buffy Coat: list of published papers using this protocol

- Boydum A. Isolation of mononuclear cells and granulocytes from human blood. Scand.J.Clin.Lab. Invest. 21 (Suppl.97): 77-89, 1968

- Alex Mabou Tagne, Franca Marino, Massimiliano Legnaro, Alessandra Luini, Barbara Pacchetti and Marco Cosentino. A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions. Int J Mol Sci 2019 Apr; 20(8): 1833. Published online 2019 Apr 13. doi: 10.3390/ijms20081833.

- Angela Scanzano, Laura Schembri, Emanuela Rasini, Alessandra Luini, Jessica Dallatorre, Massimiliano Legnaro, Raffaella Bombelli, Terenzio Congiu, Marco Cosentino, Franca Marino. Adrenergic Modulation of Migration, CD11b and CD18 Expression, ROS and interleukin-8 Production by Human Polymorphonuclear Leukocytes. Inflamm Res. 2015 Feb;64(2):127-35. doi: 10.1007/s00011-014-0791-8. Epub 2015 Jan 6.

MATERIALS

- Ethylenediaminetetraacetic acid disodium salt dihydrate: Sigma AldrichCatalog #ED2SS
- Ficoll Paque PLUS: Ge HealthcareCatalog #17144003-500 ml
- Fetal Bovine Serum (FBS): EuroCloneCatalog #ECS0180L-500 ml
- RPMI 1640: EuroCloneCatalog #ECM 0495L- 500 ml
- NaCl: Sigma AldrichCatalog #S9625
- NH4Cl: Merck Serono GmbHCatalog #1.01145.1000
- KHCO3: Merck Serono GmbHCatalog #1.04854.500
- Acetic Acid 100%: Sigma AldrichCatalog #A6283
- Genitian violet 1%: Marco VitiCatalog #not available

Optical Microscope

BEFORE START INSTRUCTIONS

All reagents used in this protocol must be at room temperature

- 1 Place 5 mL of venus blood from BUFFY COAT into 10 ml volume centrifuge tube.
- 2 Add 2 mL of **SOLUTION- 03** and mix well drawing in and out of a pipette.

Document

NAME

SOLUTION- 03 - Dextran solution 5%

CREATED BY

Farmacologia Medica

PREVIEW

- 3 Incubate in the **DARK** for 00:45:00 at 37 °C

45m

4 Place 3 mL of Fycoll-HyPaque media solution into a 10 ml volume centrifuge tube.

5 Slowly and carefully layer the supernatant from blood/dextran mixture onto the Fycoll-HyPaque media solution.

Note

Important: when layering the sample, do not mix the Fycoll-HyPaque media solution and supernatant

6 Centrifuge at 400 x g, Room temperature, 00:30:00 , no break

Equipment

Allegra AVANTI 30

NAME

Centrifuge

TYPE

Beckman Coulter

BRAND

Beckman Italy

SKU

7 Draw off the mononuclear cell layer at the Ficoll/plasma interface along with plasma and Ficoll media, leaving the white cell layer of granulocytes above the red blood cell layer undisturbed.

- 8 Resuspend the remaining cell layer in 5 mL of NaCl 0.15 Molarity (M) (**SOLUTION- 01**) and centrifuge at 400 x g, Room temperature, 00:05:00

Document

NAME

SOLUTION- 01 - Sodium Chloride (NaCl) solution

CREATED BY

Farmacologia Medica

PREVIEW

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

9 Aspirate the supernatant with a plastic Pasteur pipette.

10 Lyse remaining red blood cells in 5 mL of hypotonic Lysis Buffer (**SOLUTION- 06**) for 00:05:00

5m

Document
NAME
	SOLUTION- 06 - Lysis Buffer
CREATED BY	
Elisa Storelli	PREVIEW

11 Centrifuge at 400 x g, Room temperature, 00:05:00

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

12 Aspirate the supernatant with a plastic Pasteur pipette.

13 Resuspend the pellet in 5 mL NaCl 0.15 Molarity (M) (**SOLUTION- 01**)

Document
NAME
	SOLUTION- 01 - Sodium Chloride (NaCl) solution
CREATED BY	
Farmacologia Medica	<u>PREVIEW</u>

14 Centrifuge at 400 x g, Room temperature, 00:05:00

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

15 Aspirate the supernatant with a plastic Pasteur pipette.

16 Resuspend the cell pellet in 5 mL NaCl 0.15 Molarity (M) (**SOLUTION- 01**) for manual cell counting with Türk solution (follow protocol CELL COUNT- 02).

Document
NAME
	SOLUTION- 01 - Sodium Chloride (NaCl) solution
CREATED BY	
Farmacologia Medica	PREVIEW

Document
NAME
	SOLUTION- 08 - Türk solution
CREATED BY	
Farmacologia Medica	PREVIEW

17 OPTIONAL STEP

Cell viability can be checked using protocol CELL COUNT- 03 in the automated cell counter with Trypan Blue solution.

Document

NAME

SOLUTION- 09 - Trypan Blue solution

CREATED BY

Farmacologia Medica

PREVIEW

Equipment

Cellometer Auto T4

NAME

Automated Cell Counter

TYPE

Nexcelom Bioscience

BRAND

Euroclone

SKU

18 OPTIONAL STEP

If needed, check the purity of PMN suspension by using morphological parameters of the flow cytometer.

For this test $0,5 \times 10^6$ PMN in 500 μL of PBS (**SOLUTION- 02**) are enough.

Document

NAME

SOLUTION- 02 - Phosphate Buffered Saline (PBS)

CREATED BY

Farmacologia Medica

PREVIEW

Equipment

BD FACS Celesta

NAME

Flow Cytometer

TYPE

Becton Dickinson

BRAND

Milan Italy BD

SKU

19 EXPECTED RESULTS

Expected result

VIABILITY: the expected viability by Tripan Blue should be $\geq 90\%$

CELL YIELD: $\pm 6 \times 10^6$ cells starting from 1 mL of Buffy Coat