VERSION 2

NOV 21, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bpxzmpp6

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PMN- 02 - MTT Assay in Human PMN. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bpxzmpp6> Version created by **Farmacologia Medica**

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 21, 2020

Last Modified: Nov 21, 2020

PROTOCOL integer ID:
 44761

PMN- 02 - MTT Assay in Human PMN V.2

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Farmacologia Medica

ABSTRACT

Published work using this protocol:

- A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions.
 Mabou Tagne A, Marino F, Legnaro M, Luini A, Pacchetti B, Cosentino M. Int J Mol Sci. 2019 Apr 13;20(8):1833. doi: **10.3390/ijms20081833**

MATERIALS

MATERIALS

- ☒ Dimethyl sulfoxide (DMSO) **Sigma Aldrich Catalog #276855**
- ☒ Fetal Bovine Serum (FBS) **EuroClone Catalog #ECS0180L-500 ml**
- ☒ RPMI 1640 **EuroClone Catalog #ECM 0495L- 500 ml**
- ☒ Thiazolyl blue tetrazolium bromide (MTT) **Sigma Aldrich Catalog #M5655**
- ☒ Penicillin/Streptomycin **EuroClone Catalog #ECB3001D - 100 ml**

Instrumentation needed:

Sterile and non sterile plastic disposables
 ELISA plate reader

BEFORE START INSTRUCTIONS

Use all reagents at Room Temperature

- 1 Isolate PMN according to the protocol PMN-01a for buffy coats or PMN-01b for fresh whole blood.
- 2 Resuspend PMN at 1×10^6 cells/ml in RPMI 1640 medium supplemented with 10% Fetal Bovine Serum (FBS) and 1% penicillin/streptomycin (**SOLUTION- 13**).

Document

NAME

SOLUTION- 13 - Complete culture medium

CREATED BY

Farmacologia Medica

PREVIEW

- 3 Add 250 μ L of cell suspension per well in a 96-well round bottom plate and incubate for 24:00:00 alone or in the presence of the test substance at 37 °C in a 5% CO₂ atmosphere. **Cells must be plated in duplicate for each experimental condition.**
- 4 Centrifuge at 1400 x g, 00:05:00 and discard the supernatant.

Equipment

Allegra AVANTI 30

NAME

Centrifuge

TYPE

Beckman Coulter

BRAND

Beckman Italy

SKU

- 5 Resuspend cells with 200 μ L of **MTT solution at 0.5 mg/ml**, incubate for 02:00:00 at

37 °C in a 5% CO₂ atmosphere.

6 Centrifuge at 1400 x g, 00:05:00 and discard the supernatant.

Equipment	
Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

7 Add 120 µL of **DMSO** per well, mix the plate until the formazan crystals are dissolved.

8 Read the plate on the ELISA plate reader at 570 nm.