JUL 07, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bhrmj546

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PMN- 03 Culture of Human PMN - Migration. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bhrmj546>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 22, 2020

Last Modified: Jul 07, 2020

PROTOCOL integer ID:
 38413

PMN- 03 Culture of Human PMN - Migration

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Mattia Di Rocco

ABSTRACT

Published work using this protocol:

- A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions.

Alex Mabou Tagne, Franca Marino, Massimiliano Legnaro, Alessandra Luini, Barbara Pacchetti, Marco Cosentino. Int J Mol Sci. 2019 Apr 13;20(8):1833. doi: 10.3390/ijms20081833.

- β 2 -Adrenoceptors Inhibit Neutrophil Extracellular Traps in Human Polymorphonuclear Leukocytes.

Franca Marino, Angela Scanzano, Laura Pulze, Monica Pinoli, Emanuela Rasini, Alessandra Luini, Raffaella Bombelli, Massimiliano Legnaro, Magda de Eguileor, Marco Cosentino . J Leukoc Biol. 2018 Sep;104(3):603-614. doi: 10.1002/JLB.3A1017-398RR. Epub 2018 Apr 18.

- Adrenergic Modulation of Migration, CD11b and CD18 Expression, ROS and interleukin-8 Production by Human Polymorphonuclear Leukocytes.

Angela Scanzano, Laura Schembri, Emanuela Rasini, Alessandra Luini, Jessica Dallatorre, Massimiliano Legnaro, Raffaella Bombelli, Terenzio Congiu, Marco Cosentino, Franca Marino. Inflamm Res. 2015 Feb;64(2):127-35. doi: 10.1007/s00011-014-0791-8. Epub 2015 Jan 6.

MATERIALS

MATERIALS

⊗ RPMI 1640 EuroClone Catalog #ECM 0495L- 500 ml

⊗ N-FORMYL-MET-LEU-PHE (fMLP) Merck MilliporeSigma (Sigma-Aldrich) Catalog #F3506

⊗ Propanolol Merck MilliporeSigma (Sigma-Aldrich) Catalog #P0884

⊗ Xylene Merck MilliporeSigma (Sigma-Aldrich) Catalog #247642

⊗ Hematoxylin Merck MilliporeSigma (Sigma-Aldrich) Catalog #MHS16

⊗ IL-8 Merck MilliporeSigma (Sigma-Aldrich) Catalog #I1645

STEP MATERIALS

⊗ RPMI 1640 EuroClone Catalog #ECM 0495L- 500 ml

⊗ IL-8 Merck MilliporeSigma (Sigma-Aldrich) Catalog #I1645

⊗ N-FORMYL-MET-LEU-PHE (fMLP) Merck MilliporeSigma (Sigma-Aldrich) Catalog #F3506

⊗ Propanolol Merck MilliporeSigma (Sigma-Aldrich) Catalog #P0884

⊗ Hematoxylin Merck MilliporeSigma (Sigma-Aldrich) Catalog #MHS16

⊗ Xylene Merck MilliporeSigma (Sigma-Aldrich) Catalog #247642

Dubnoff bath

1 Separation of **PMN** (at least 7 million PMN).

2 Preparation of 6 Boyden chambers:

Cut the bottom part of the vial and apply the filter (with holes of 3-5 μm in diameter) with glue / mastic (fig. 1). With paper scotch, name all Boyden chambers (fig. 2).

figure 1

figure 2

- 3 Preparation of the solutions for:
 - **IL-8** 10 ng/ml (2 μ L stock in 198 μ L di RPMI) chemotactic agent,
 - **fMLP** 0.1 μ M (10 μ L stock in 990 μ L di PBS/BSA) chemotactic agent,
 - other activating stimulus according specific experimental plan,
 - drug/substances to test at different concentrations: one chamber for each concentration.

⊗ **IL-8 Merck MilliporeSigma (Sigma-Aldrich) Catalog #I1645**

⊗ **N-FORMYL-MET-LEU-PHE (fMLP) Merck MilliporeSigma (Sigma-Aldrich) Catalog #F3506**

- 4 Fill the bottom of each Boyden chamber with 1 mL of **RPMI**.

- 5 Remove 10 μL of **RPMI** from each Boyden chamber and add 10 μL of **IL-8** or **fMLP** treatment (except for control).
- 6 Resuspend 7×10^6 PMN in 700 μL of **RPMI**.
- 7 Immerse the filters in the chambers.
- 8 Put 100 μL of cell suspension in 4 vials and treat cells with 10 μL of testing drug.
- 9 Place the cells on the filters (100 μL of cell suspension on each filter) using the P100 taking care to direct the tip towards the center of the filter without touching it (if the filter does not get wet at all, gently slam the chamber).
- 10 Cap and put Boyden's chambers in the Dubnoff bath for 01:30:00 .
- 11 Under the chemical hood, prepare the following material:
 - 4 rows of mini becker (6 becker per row).
In each row of becker arrange, in the following order, equal quantities (2 ml in total) of: **water**, **propanolol**, **xylene**, **propanolol + xylene** (fig. 3)
 - 1 row of lids for 3-6 drops of hematoxylin to cover the same area of the filter (6 lids) and 5 plastic pasteur (each suitably marked with the name of the substance that will collect: **water**, **propanolol**, **xylene**, **hematoxylin**, immersion oil).(fig.4)

Figure 3

Figure 4

12 Immerse the filters related to eppendorf in:

■ **PROPANOLOL** for 00:04:00

■ **WATER** for 00:04:00

■ **HEMATOXYLIN** for 00:06:00

■ **WATER** for 00:06:00

(in the meantime throw the hematoxylin into the liquid discharge, wrap its containers in the paper and throw it all in special waste)

■ **PROPANOLOL** for 00:06:00

■ **PROPANOLOL + XYLENE** for 00:06:00

(meanwhile throw away water and propanolol in the liquid discharge and immerse the beakers to be washed in a bowl with water, alcohol and bleach taking care to turn them over so they are hygienized every side)

XYLENE for 00:06:00

⊗ Propanolol Merck MilliporeSigma (Sigma-Aldrich) Catalog #P0884

⊗ Xylene Merck MilliporeSigma (Sigma-Aldrich) Catalog #247642

⊗ Hematoxylin Merck MilliporeSigma (Sigma-Aldrich) Catalog #MHS16

13 Throw away the used pasteur, wrapping it first in the paper.

14 Set up the slides:
nominate them,
arrange the filter (taking care to maintain the correct orientation),
arrange the cover glass (by tapping laterally with the tweezers to make sure it is well stuck),
add on top a drop of immersion oil.

15 Microscopy: make sure that the lens comes into contact with the immersion oil.

16 Once the filter with the colored cells in Haematoxylin is in focus, using only the micrometric lens bring into focus the layer of neutrophils above and report the value by saying "UP"; then you go down into the depth of the filter (rotating the micrometric lens) and bring into focus the migrated neutrophil layer: mark the visual limit value of the individual migrated cells saying "DOWN".

17 The delta between the "UP" and "DOWN" values (express as μm) represents the distance in

micrometers traveled by the migrating cells.

- 18** For each slide make 10 counts of the distance traveled by migrated neutrophils (selected in different optical fields): each count must be done by focusing on a different migrated neutrophil.
- 19** Results will be expressed or as absolute migration toward the filter or as ratio vs activating stimulus (sample migration/stimulus migration)