VERSION 2

NOV 21, 2020

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.bpx2mpqe

Protocol Citation: Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. PMN- 06 - Culture of Human PMN - TNF- α production.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bpx2mpqe> Version created by Farmacologia Medica

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Nov 21, 2020

Last Modified: Nov 21, 2020

PMN- 06 - Culture of Human PMN - TNF- α production V.2

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹,
Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹,
Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese, Italy)

Farmacologia Medica

ABSTRACT

Published work using this protocol:

- A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions. Mabou Tagne A, Marino F, Legnaro M, Luini A, Pacchetti B, Cosentino M. Int J Mol Sci. 2019 Apr 13;20(8):1833. doi: 10.3390/ijms20081833

MATERIALS

MATERIALS

Fetal Bovine Serum (FBS) EuroClone Catalog #ECS0180L-500 ml

RPMI 1640 EuroClone Catalog #ECM 0495L- 500 ml

Lipopolysaccharides from Escherichia coli O127:B8 Sigma Catalog #L3137

Penicillin/Streptomycin EuroClone Catalog #ECB3001D - 100 ml

ELISA kit for TNF- α Thermo Fisher Scientific Catalog #KHC0031

Cannabidiol Linnea SA Catalog #not available

Instrumentation needed:

Sterile and non sterile plastic disposables
ELISA plate reader

- 1 Isolate PMN according to the protocol PMN-01a or PMN-01b.
- 2 Resuspend PMN at 10×10^6 cells/ml in RPMI medium supplemented with 10% FBS and 1% penicillin/streptomycin (**SOLUTION- 13**).

Document

NAME
SOLUTION- 13 - Complete culture medium

CREATED BY
Farmacologia Medica

PREVIEW

- 3 Add 300 μ L of cell suspension in a 5 ml tube for cell culture and incubate for 21:00:00 at 37 °C in 5% CO₂. Cells were incubated in resting condition or in presence of stimulus. Stimulus (**LPS 1 μ g/ml** and/or **CBD 10⁻⁸-10⁻⁵M**) in 5 ml tubes for cell culture. **Other possible activating stimuli** can be selected and included.

- 4 Centrifuge at 1500 x g, 00:10:00

Equipment

Allegra AVANTI 30	NAME
Centrifuge	TYPE
Beckman Coulter	BRAND
Beckman Italy	SKU

- 5 Harvest cell-free supernatant (150 µL x2 aliquots) and cell pellet.
- 6 Store the supernatant at -80°C until the TNF-α Enzyme Linked-Immuno-Sorbent Assay (ELISA) is performed.
- 7 Store cell pellet at -80°C for gene expression analysis (RT PCR).