VERSION 2

AUG 16, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.sn8edhw

Protocol Citation: Tomasz Suchan 2018. Pollen metabarcoding. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.sn8edhw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
correcting some typos

Created: Aug 16, 2018

Last Modified: Aug 16, 2018

PROTOCOL integer ID:
14784

Pollen metabarcoding V.2

Tomasz Suchan¹

¹Institute of Botany, Polish Academy of Sciences

Molecular Biogeography Group

Tomasz Suchan

W. Szafer Institute of Botany, Polish Academy of Sciences

GUIDELINES

Perform reactions in small batches until you are confident that there is no cross-contamination among the samples. Including isolation blanks and PCR blanks is crucial for the quality control.

MATERIALS

MATERIALS

Q5 Hot Start High-Fidelity DNA Polymerase - 100 units **New England Biolabs Catalog #M0493S**

Water, nuclease free **Contributed by users**

dNTP mix (25 mM of each) **Contributed by users**

Phire Plant Direct PCR Kit **Thermo Fisher Scientific Catalog #F130WH**

BEFORE START INSTRUCTIONS

Prepare 5 μ M primer solutions:

ITS2 primers used in the 1st PCR:

ITS2-4R

GTGACTGGAGTTCAGACGTGTGCTCTTCCGATCTNNNNNNTCCTCCGCTTATTGATA
TGC

ITS2-S2F

ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNNNATGCGATACTTGGTGTGA
AT

Indexing primers used in the 2nd PCR (xxxxxxx - index):

AATGATACGGCGACCACCGAGATCTACACxxxxxxxACACTCTTTCCCTACACGACGC
CAAGCAGAAGACGGCATACGAGATxxxxxxxGTGACTGGAGTTCAGACGTGTGC

Pollen extraction

- 1 Vortex the butterfly in 50 μ l of water with 0.1% SDS.

Safety information

Add blank sample at this step (= "isolation blank")

<http://null>

- 2 Evaporate water in speedvac.
- 3 Add 5 μ l of the Phire Plant Direct sample buffer.
- 4 Spin max speed for 2 min.

1st PCR

5 Prepare the mix:

14 µL molecular grade water

25 µL Phire Plant Direct PCR mix

5 µL ITS2-S2F primer

5 µL ITS2-4R primer

6 Add 1 µl of the sample to 49 µl of the mix. Use water instead of the sample for the blanks.

Safety information

Add another blank sample at this step (= "PCR blank")

7 Run the PCR program: initial denaturation at 98°C for 5 min; 20 cycles of denaturation at 98°C for 40 s, annealing at 49°C for 40 s and elongation at 72°C for 40 s; followed by a final extension step at 72°C for 5 min.

Purification

8 Perform AMPure purification with ratio 1x. Elute in 10 µl.

Protocol

NAME

AMPure purification protocol

CREATED BY

Tomasz Suchan

PREVIEW

8.1 Remove beads from the fridge, equilibrate to the room temperature, mix thoroughly but do

not vortex.

- 8.2** Add desired ratio of AMPure beads to the DNA to be purified and mix well by pipetting.
- 8.3** Incubate 5 minutes.
- 8.4** Place on the magnetic rack.
- 8.5** Let it stand for 5 minutes on the rack, aspirate and discard supernatant.
- 8.6** Add 200 µl of freshly prepared 70% EtOH, incubate 30 seconds, aspirate and discard EtOH.
- 8.7** Repeat the wash: add 200 µl of freshly prepared 70% EtOH, incubate 30 seconds, aspirate and discard EtOH.
- 8.8** Wait until the pellet dries out completely. DO NOT remove the plate/tubes from the magnetic rack, as the dried AMPure might pop out from the tubes!
- 8.9** Add desired volume of Tris 10 mM or water, wetting the dried AMPure pellets (add 1 µl to the final volume to avoid pipetting out the beads)

- 8.10** Remove from the magnetic rack.
- 8.11** Resuspend by pipetting or vortexing.
- 8.12** Incubate 10 minutes, incubating in 37°C can improve DNA yield.
- 8.13** Place on the magnetic rack.
- 8.14** Let it stand for 5 minutes, pipette out and save supernatant. The eluted DNA is in the supernatant, do not discard it!

2nd PCR

9 Prepare the mix:

4.82 µL molecular grade water

2 µL Q5 reaction buffer

0.08 µL dNTPs (25 mM each)

0.1 µL Q5 Hot Start polymerase

10 Add 1 µl of the template to 7 µl of the mix.

- 11** Add 2 µl of each 5 uM primer (forward and reverse).

Safety information

Use different pair of indexed primers for each sample.

- 12** Run the PCR program: 30 s denaturation at 98°C; 12 cycles of denaturation at 98 °C for 10 s, combined annealing and extension at 72°C for 30 s (shuttle PCR); the final extension at 72°C for 5 min.

- 13** Check profiles on a gel.

Expected result

Pooling

- 14** Pool all the samples together.

15 Perform AMPure purification with ratio 1x. Elute in 100 µl.

Protocol

NAME

AMPure purification protocol

CREATED BY

Tomasz Suchan

PREVIEW

- 15.1** Remove beads from the fridge, equilibrate to the room temperature, mix thoroughly but do not vortex.
- 15.2** Add desired ratio of AMPure beads to the DNA to be purified and mix well by pipetting.
- 15.3** Incubate 5 minutes.
- 15.4** Place on the magnetic rack.
- 15.5** Let it stand for 5 minutes on the rack, aspirate and discard supernatant.
- 15.6** Add 200 µl of freshly prepared 70% EtOH, incubate 30 seconds, aspirate and discard EtOH.
- 15.7** Repeat the wash: add 200 µl of freshly prepared 70% EtOH, incubate 30 seconds, aspirate

and discard EtOH.

- 15.8** Wait until the pellet dries out completely. DO NOT remove the plate/tubes from the magnetic rack, as the dried AMPure might pop out from the tubes!
- 15.9** Add desired volume of Tris 10 mM or water, wetting the dried AMPure pellets (add 1 µl to the final volume to avoid pipetting out the beads)
- 15.10** Remove from the magnetic rack.
- 15.11** Resuspend by pipetting or vortexing.
- 15.12** Incubate 10 minutes, incubating in 37°C can improve DNA yield.
- 15.13** Place on the magnetic rack.
- 15.14** Let it stand for 5 minutes, pipette out and save supernatant. The eluted DNA is in the supernatant, do not discard it!
- 16** Check the concentration using Qubit and the profile using Tapestation/Fragment Analyzer. Calculate molarity from that and proceed to the sequencing. Add 15% PhiX to the sequencing run.