AUG 10, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bjiykkfw

Protocol Citation: Angel A Justiz-Vaillant 2020. Polson method for IgY isolation with modification. . **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bjiykkfw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 10, 2020

Last Modified: Aug 10, 2020

PROTOCOL integer ID:
 40248

🌐 Polson method for IgY isolation with modification.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

ABSTRACT

Immunoglobulin Y (IgY) is the mayormolecule present in the avian egg yolk. This protein functions as antibody and fulfils important functions in the protection of the embryo against challenging stimuli [1]. IgY has been isolated from the egg yolk of many birds [1]. These antibodies are important therapeutically, but also they have been used as immunological tool in many immunodiagnostic assays.

Much time IgY concentration is calculated from the water soluble fraction (WSF), which means that there is not need of treating the preparation with alcohol to separate the protein pellet.

Reference

1. Justiz Vaillant AA, Ramirez N, Cadiz A, Ferrer B, Akpaka P, et al. (2013) Separation and Reactivity of Avian Immunoglobulin Y. J Chromat Separation Techniq 4: 173. doi:10.4172/2157-7064.1000173

GUIDELINES

The modification in the Polson method (1990) is based on the substitution of the protein separation step with PEG by another suitable reagent as ethanol or a substance as trichloroacetic acid [1].

Reference

1. Asemota H, Curtello S, Vaillant AAJ, Mohammed W, Vuma S, et al. (2013) Purification of Avian IgY with Trichloroacetic Acid (TCA). J Chromatograph Separat Techniq 4: 205. doi:10.4172/2157-7064.1000205

MATERIALS

MATERIALS

⊗ 100 ml Polyethyleneglycol 600 **biorbyt Catalog #orb65596**

⊗ Chloroform **Sigma Catalog #366919-1L**

⊗ Trichloroacetic Acid **Sigma Aldrich Catalog #490-10**

⊗ Ethanol **Merck Millipore Catalog #100983**

⊗ PBS **Life Technologies**

- 1 Wash the egg with warm water and crack the eggshell carefully.
- 2 Separate the egg yolk from the egg white manually and place the egg content on a tissue paper to help to remove as much egg white as possible.
- 3 Wash four times the egg yolk with 100 ml of PBS, pH 7.4.
- 4 Break the egg yolk membrane and pour it into a 50 ml tube. Then, dilute egg yolk 1:5 in PBS, pH 7.4, and add an equal volume of chloroform (ACS reagent grade, Sigma-Aldrich).
- 5 Shake the mixture, and vortex and roll the tube containing the mixture on rolling mixer for 15 minutes.
- 6 Decant the supernatant, which contains the water-soluble fraction (WSF) that is the diluted egg yolk plasma, rich in IgY (25 mg/ml), which is approximately four times the IgY concentration present in the serum (6 mg/ml).
- 7 Precipitate egg yolk antibodies (IgY) by dropping the WSF with cold ethanol, trichloroacetic acid (TCA) or the classic polyethylene glycol. (PEG).

- 8 Estimate the protein concentration.
- 9 Run a protein gel electrophoresis (PAGE) of the IgY to assess the molecular weight of the chicken egg yolk antibody, protocol for this can be found elsewhere.
- 10 Then use Phosphate-buffered saline (PBS) pH=7.4 to dialyze the preparations for 8 hrs at 4°C.
- 11 The dialysate is stored at -20°C until further use.
- 12
- 13