NOV 26, 2018

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.vwae7ae

Protocol Citation: Karen Guimaraes 2018. Polymerase Chain Reaction (PCR) - DNA barcoding. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.vwae7ae>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Nov 26, 2018

Last Modified: Nov 26, 2018

PROTOCOL integer ID:
 18082

Polymerase Chain Reaction (PCR) - DNA barcoding

Karen Guimaraes¹

¹Laboratório de Genética & Biodiversidade, Instituto de Ciências da Educação, Universidade Federal do Oeste do Pará, Santarém, Pará, Brazil

Karen Guimaraes

Laboratório de Genética & Biodiversidade, Instituto de Ciênc...

GUIDELINES

PCR conditions: 95°C (2min), 35 cycles of 94°C (30sec), 54°C (30sec) and 72°C (1min), followed by 72°C (10min).

BEFORE START

The reactions in 25 µL final volume.

1. Add 15 µL sterile H₂O, 2.8 µL dNTP mix (1.25 mM);
2. Add 2.5 µL buffer 10X (200 mM Tris-HCl (pH = 8,4) + 500 mM KCl);
3. Add 2.5 µL de MgCl₂ (50 mM);
4. Add 0.5 µL of each primer (5µM);
5. Add 0.2 µL Taq DNA polymerase (5U/µL) and 1 µL of genomic DNA (100ng/µL)

MATERIALS

Sterile H₂O; dNTP mix (1.25 mM); buffer 10X (200 mM Tris-HCl (pH = 8,4) + 500 mM KCl); MgCl₂ (50 mM); Primer Fish F1 and Fish R1 (5µM); Taq DNA polymerase.

BEFORE START INSTRUCTIONS

The reactions in 25 µL final volume.

1. Add 15 µL sterile H₂O, 2.8 µL dNTP mix (1.25 mM);
2. Add 2.5 µL buffer 10X (200 mM Tris-HCl (pH = 8,4) + 500 mM KCl);
3. Add 2.5 µL de MgCl₂ (50 mM);
4. Add 0.5 µL of each primer (5µM);
5. Add 0.2 µL Taq DNA polymerase (5U/µL) and 1 µL of genomic DNA (100ng/µL)