FEB 28, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.ba79ihr6

External link:
<https://www.addgene.org/protocols/pooled-library-amplification/>

Protocol Citation: Addgene The Nonprofit Plasmid Repository 2020. Pooled Library Amplification.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.ba79ihr6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jan 10, 2020

Last Modified: Feb 28, 2020

PROTOCOL integer ID:
 31713

Pooled Library Amplification

Addgene The Nonprofit Plasmid Repository¹

¹Addgene

Addgene The Nonprofit Plasmid Repository
 Addgene

ABSTRACT

This protocol describes pooled library amplification. To see the full abstract and additional resources, visit

<https://www.addgene.org/protocols/pooled-library-amplification/>.

GUIDELINES

Required Quality Control (QC):

Pooled libraries can be challenging and expensive to ensure adequate quality, but this upfront cost will save headaches and expense later.

At a minimum, we recommend the use of a diagnostic digest and high-throughput next-generation sequencing (NGS). Select a restriction enzyme for digest that will cut the shared plasmid backbone a single time and visualize that digestion on an agarose gel ([see protocol here](#)). Lentiviral plasmids can recombine between their LTRs resulting in a smaller plasmid containing only the elements required for bacterial propagation (origin of replication and antibiotic selection). This recombination, at a low rate, is not typically a problem as these sequences are not efficiently packaged into lentiviral particles. See below for options if the recombinant band makes up a significant proportion of the DNA pool.

Workflow Timeline:

Day 1: Transform, recover, set up overnight growth (Estimated time 02:00:00

03:00:00

Note

Transformation should be performed at the end of the day to ensure that growth times are limited.

Keywords: Pooled library, amplification, pooled plasmid library, screen

Day 2: Harvest cells and purify DNA (Estimated time 03:00:00 - 04:00:00)

Note

Cells should be harvested first thing in the morning to ensure that outgrowth and competition are limited.

Tips and Troubleshooting

What do I do if my transformation efficiency is not high enough?

Repeat the experiment with more cells, more DNA, gentler practices, and colder reagents (not including the recovery media!). Do not proceed with Maxipreps or NGS until adequate transformation efficiency is obtained. Ensure that electrocompetent cells are being used. Chemically competent cells will not provide adequate transformation efficiency! Ensure that no arcing is taking place during electroporation. Arcing would manifest as a loud pop often accompanied by some light in the cuvette during the electroporation step.

I got a heavy recombined band by digest

Recombination is thought to be in part related to selective pressure. These recombined plasmids are much smaller and are a smaller nutritional burden on the bacteria. Bacteria that contain a recombinant plasmid likely grow faster in the closed system of a crowded plate or culture. Try restricting growth to the bare minimum by growing at 30°C, limiting growth time further, or potentially spreading less bacteria on a given plate (ex by increasing the total number of bioassay plates used). Scaling up the experiment using a Gigaprep can be used to obtain sufficient intact pooled library even in the face of a recombined fraction but care must be taken to perform adequate NGS based QC to ensure no change in representation compared to the pre-amplified stock. A last option is to perform a [gel extraction](#) of either the original sample or the amplified sample, followed by reamplification of the DNA, but please note that NGS should be performed to ensure representation is maintained.

Maxipreps - Less is more

Do not overload the Maxipreps as yield can dramatically plateau and sometimes fall off entirely if the column or reagents are significantly overloaded. If you find that your pellet outstrips the capacity listed in the Maxiprep protocol, scale the reagent volume and column number as needed. The use of Mega or Gigapreps is acceptable when scaling up or reducing the number of tandem purifications. Consult the manufacturers' handbooks for appropriate volumes and numbers of tandem purifications.

MATERIALS

Equipment

- Table top centrifuge
- BioRad Electroporator (MicroPulser TM , Bio-Rad 1652100)

Reagents

- 200 µL electrocompetent cells (Default: 4 tubes of Endura Duos, Lucigen, 60242-1)

Note

- Alternatives include Stbl4 cells or other ultra-high efficiency electrocompetent cells that are suitable for unstable or recombination-prone DNA.
- The use of electrocompetent cells is essential to ensure high efficiency uptake of plasmid library DNA.
- This quantity of cells is sufficient for libraries up to 200,000 individual plasmids. For larger libraries scale the number of cells and DNA accordingly.

Note

Pro-tip

Use of extremely high efficiency (1×10^{10} cfu/µg) commercially prepared electrocompetent cells is strongly recommended.

- 800 ng pooled library DNA (100 ng plasmid DNA per 25 µL electrocompetent cells)

Note

Less library DNA can be used but this dramatically increases the chances of individual plasmids being lost from the pool and/or increasing the skewness of the pool.

- 8 electroporation cuvettes (BioRad, Micropulser, 0.1 cm)
- 20 mL SOC recovery media (Lucigen, 80026-1)
- 8X LB Agar + Antibiotic 245 mm bioassay plates (Molecular Devices, X6023)

Note

Pro-tip

Pour these plates at least one day in advance to allow adequate time to fully gel and to dry slightly. We routinely use 350 mL of LB Agar per bioassay plate.

- 14 mL Vented Falcon Tubes (BD Biosciences, 352059)
- 3X LB Agar + Antibiotic 65 mm Petri dish (VWR, 11019-552)
- 4 Maxipreps (Qiagen HiSpeed Max, Catalog #12663)
- Tips (1000 µL, 200 µL, 10 µL)
- Bacti Cell Spreaders (VWR, 60828-680)
- 5 mL and 10 mL Serological pipettes
- Ice slurry (Ice bucket with ice and water to create slurry)
- 100 mL LB house-made

Note

At Addgene we use premixed LB (VWR 101414-072) for convenience but any brand or house-made LB that supports normal growth is expected to work

- 50 mL Falcon Conical tubes (Fisher, 14-432-22)

Reagent Preparation (optional)

1. Prepare, sterilize, and pour all LB Agar + Antibiotic plates.
2. Prewarm 12 mL recovery media at 37 °C (for at least 00:15:00).
3. Prewarm 3X LB Agar + Antibiotic plates at 37 °C.
4. Prewarm 8X LB Agar + Antibiotic Bioassay plates.
5. Prechill Micropulser cuvettes on ice.
6. Thaw 4 tubes of electrocompetent cells on ice for 00:15:00 - 00:20:00 or until completely thawed.
7. Chill a box of 200 µL micropipette tips in a -20 °C freezer.
8. Aliquot 3 mL SOC into each of four 14 mL Vented Falcon Tubes and have 1 mL SOC per electroporation readily available for post-electroporation recovery of cells.
9. Ensure access to autoclaved, sterile reagents for all steps.

SAFETY WARNINGS

See SDS of listed reagents for safety warnings and hazards.

Day 1

1 Add 200 ng DNA to each 50 μ L aliquot of thawed Endura Duos on ice. Flick gently to mix

2 Electroporate cells (one at a time for a total of eight electroporations):

Note

Electroporator Conditions: Bio-Rad Micropulser Ec1 0.1 cm cuvette, 1.8 kV, 1 pulse.

3 Aliquot 25 μ L DNA-Endura into pre-chilled cuvette.

4 Pulse.

Safety information

Electroporation involves the use of high voltages, please use caution when activating pulse and follow all specifications described in the equipment manual.

5 Immediately add 1 mL SOC to cuvette.

6 Remove all liquid from cuvette and add to 14 mL vented Falcon Tube containing 3 mL SOC.

- 7 Repeat for each of the 25 μL aliquots of cell/DNA mixture used (8 in the described protocol) using a new vented falcon tube for every two transformations. Each of the four 14 mL vented falcon tubes should contain a total of 5 mL (3 mL SOC + 2 mL transformed Endura from two separate transformations).

Note

Pro-Tip

Do not pipette repeatedly or mix when removing SOC containing transformed DNA-Endura from cuvette.

- 8 Shake four 14 mL Vented Falcon Tubes at 30 °C 37 °C , 225 rpm for 01:00:00 .
- 9 After the 1 hour shaking period, pool and gently mix the four tubes.
- 10 Perform sequential 1:100 dilutions of the cells (add 10 μL of the pool to 990 μL LB then perform a second and then third 1:100 dilution). Plate 100 μL of each dilution onto a prewarmed Petri dish.
- 11 Incubate plates at 30 °C overnight.
- 12 Plate 2.5 mL of the transformed cells on each of the eight bioassay plates (two plates per tube). Distribute evenly with a sterile spreader until all liquid has been absorbed by the agar. This usually takes 00:01:00 00:02:00 .

Note

Critical

Be careful not to rip or shred the agar. Do so by gentle spreading. Some spreaders have a sharp edge that can scrape plates more abrasively at a certain angle.

- 13 Incubate plates upside down at 30 °C overnight.

Note

Critical

Ensure at this stage that no unabsorbed media drips onto the lid. Let plates remain agar side up until dried before overnight incubation if needed.

- 14 Place 100 mL sterile LB at 4 °C .

Day 2

- 15 Before beginning, prechill at least four conical tubes on ice and ensure access to sterile scraper and cold LB.

Note

Pro-Tip

Prepare one to two more conical tubes on ice in case you need to spread out the harvested cells further than four Maxipreps worth.

- 16 Count colonies on the most dilute Petri dish.

Note

- Total colony yield = count x 100 x 100 x 100 ÷ 0.1.
- This number should be at least 1000x greater than the number of perturbations in the library.
- *Example: At least 10 colonies on the most diluted plate for 1000x coverage of a library of 100,000 plasmids.*
- Colonies may appear small and require extra incubation time in order to be enumerated accurately.
- Frequently the number of colonies can be too great to count. Ideally, dilutions would have been sufficient to enumerate single colonies but as long as one can ensure more than the required colonies are present. If colonies are not present on the most dilute plate, count the second plate (ie. the 1:10,000 dilution).

17 After 12:00:00 - 18:00:00 of growth, use the spreader and cold LB to scrape bioassay plates and remove bacteria.

18 Use one scraper for all plates.

19 Use two 10 mL pipettes (one for dispensing cold LB and one for removing LB containing bacteria from the plate).

Note

Pro-Tip

Scraped bacteria in LB can clog the pipette either mix gently up and down avoiding introducing bubbles or pour off plates into conical tubes as needed.

20 Add 10 mL cold LB to each plate for each scrape and use spreader to scrape plates.

Note

Pro-Tip

Pushing motion is better than pulling motion and take care not to split or gouge agar during the scraping process.

21 Add each scrape into a 50 mL conical tube on ice pooling the scrapings of two plates into each tube. Keep each tube on ice while scraping.

22 Repeat addition of 10 mL cold LB and scrape for each plate until agar is clear (all bacteria have been removed). This should produce 25 mL cold LB-bacteria per plate.

23 Centrifuge tubes (4000 x g, 4°C, 00:15:00) to pellet bacteria.

24 Decant LB and weigh pellet. - The total weight of each pellet should be ~ 1 g 2 g .

Note

Pro-Tip

Make sure to weigh the empty tube beforehand! If you've already gone too far, weigh an identical empty tube. It will be close enough for the purpose of a Maxiprep.

25 Purify plasmid DNA using the Qiagen HiSpeed Maxi Kit (one conical is its own Maxiprep).

Note

Critical

Do not freeze pellets for later purification. Immediately purify them! Commercial Maxipreps rely on incremental, ordered cell lysis. *E. coli* cells are subject to lysis by freeze-thaw if not suspended in cryoprotectant like Glycerol or DMSO solutions.

26 Quantify the individual Maxipreps by Nanodrop, Picogreen, or Qubit.

Note

Each different DNA measurement type can report slightly different concentrations of DNA. In our hands fluorescent dye methods tend to report lower values than Nanodrop, although most protocols have been designed with Nanodrop as the quantification method.

27 If all Maxipreps appear to contain sufficient DNA for use, pool samples, and continue with analysis.

Note

“Sufficient” DNA will vary upon intended use. Typically yields range from several hundred micrograms to a milligram of total plasmid DNA.