JUL 25, 2020

Porcine Circovirus 3 (PCV3) Complete Genome Sequence PCR

PLOS One

Chew Yee Tan¹

¹Universiti Putra Malaysia

Chew Yee Tan
Universiti Putra Malaysia

ABSTRACT

This protocol describes four separate PCR runs to amplify the complete genome sequence of Porcine circovirus 3 (PCV3) as four overlapping segments.

MATERIALS

MATERIALS

- QIAGEN DNeasy Blood and Tissue Kit, 50 rxn **Qiagen Catalog #69504**
- MyTaq™ Red Mix 2X master mix **Bioline**
- RedSafe™ nucleic acid staining solution **iNtRON Biotechnology, South Korea**
- GelPilot 100 bp Plus Ladder **Qiagen**

OPEN ACCESS

DOI:
[dx.doi.org/10.17504/protocols.io.bdd9i296](https://doi.org/10.17504/protocols.io.bdd9i296)

External link:
<https://doi.org/10.1371/journal.pone.0235832>

Protocol Citation: Chew Yee Tan 2020. Porcine Circovirus 3 (PCV3) Complete Genome Sequence PCR. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bdd9i296>

MANUSCRIPT CITATION:
Tan CY, Opaskornkul K, Thanawongnuwech R, Arshad SS, Hassan L, Ooi PT (2020) First molecular detection and complete sequence analysis of porcine circovirus type 3 (PCV3) in Peninsular Malaysia. PLoS ONE 15(7): e0235832. doi: [10.1371/journal.pone.0235832](https://doi.org/10.1371/journal.pone.0235832)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Mar 09, 2020

Last Modified: Jul 25, 2020

PROTOCOL integer ID:
33953

- 1 DNA extraction was performed using DNeasy Blood & Tissue Kit extraction kit (Qiagen, Germany) in accordance to manufacturer's instructions. Samples involved include lung, inguinal lymph node, spleen, tonsil, kidney, mesenteric lymph node, heart, liver and brain tissues.
- 2 Four separate conventional PCR runs were performed to amplify the complete nucleotide sequence of PCV3 as four overlapping segments.
- 3 Primers, their respective cycling conditions and expected PCR product band size are tabulated below:

Primer Pair	Nucleotide Sequence (5'– 3')	Product Length (bp)	PCR Cycling Condition (Temperature / Time)						Reference
			Initial Denaturation	Number of Cycle	Denaturation	Annealing	Extension	Final Extension	
P1F	CACCGTGTGAGTGGATA TAC	854	94°C / 4 min	35	94°C / 20 s	55°C / 30 s	72°C / 30 s	72°C / 5 min	Palinski R, Piñeyro P, Shang P, Yuan F, Guo R, Fang Y, et al. A novel porcine circovirus distantly related to known circoviruses is associated with porcine dermatitis and nephropathy syndrome and reproductive failure. J Virol. 2017 Jan 1;91(1):e01879-16.
P1R	CAAACCCACCTTAACA G								
P2F	GTCGTCTTGAGCCAAG TG	807	95°C / 5 min	35	94°C / 30 s	62°C / 30 s	72°C / 1 min	72°C / 7 min	
P2R	CGACCAAATCCGGGTAA GC								
KF	TTACTTAGAGAACGGAC TTGTAACG	649	94°C / 5 min	35	94°C / 30 s	55°C / 30 s	72°C / 1 min	72°C / 10 min	Ku X, Chen F, Li P, Wang Y, Yu X, Fan S, et al. Identification and genetic characterization of porcine circovirus type 3 in China. Transbound Emerg Dis. 2017 Jun;64(3):703-8.
KR	AAATGAGACACAGAGCT ATATTCAG								
V4F	GAAAACGCGGGAAGCTT GTG	806	95°C / 5 min	35	94°C / 30 s	56°C / 30 s	72°C / 1 min	72°C / 10 min	Designed in this study based on PCV3 strain KY075986
V4R	CCACTTCTGGCGGGAAC TAC								

- 4 Type, concentration and volume of reagents used in the four separate PCR runs are the same: 12.5 µL of MyTaq™ Red Mix 2X master mix (Bioline, United Kingdom) and 1.0µM of each primer in a pair, in a 25 µL total PCR reaction volume.

- 5 PCR products were stained using RedSafe™ nucleic acid staining solution (iNtRON Biotechnology, South Korea) and analysed by agarose electrophoresis using 2.5% agarose gel with GelPilot 100 bp Plus Ladder as marker (Qiagen, Germany).