MAY 14, 2018

Pour plating protocol for *Emiliana huxleyi* (single colonies)

Inbal Nussbaum¹, Daniella Schatz¹

¹Weizmann Institute of Science

Protist Research to Optimize Tools in Genetics (PROT-G)

Daniella Schatz

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.p4ydxw

Protocol Citation: Inbal Nussbaum, Daniella Schatz 2018. Pour plating protocol for *Emiliana huxleyi* (single colonies). **protocols.io** <https://dx.doi.org/10.17504/protocols.io.p4ydxw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: May 14, 2018

Last Modified: May 14, 2018

PROTOCOL integer ID:
12152

ABSTRACT

A protocol for generating single colonies of *Emiliana huxleyi* embedded in soft-agarose.

1 Autoclave 75 mL DDW with 0.75 g agarose

- 2 For 300 mL (= 7 plates): heat 225 mL FSW in a 50 °C water bath.
- 3 Cool autoclaved agar to 50 °C, mix with heated FSW, add F/2 stock and antibiotics if necessary. Swirl/mix well.
- 4 For each sample, pour 40 mL of FSW-agar mixture to a 50 mL conical tube.
- 5 Move to 32 °C bath.
- 6 Prepare the cells at desired cell concentration. For each sample prepare 0.5ml cells.
- 7 Lay out plates.
- 8 When you are ready, remove FSW-agar mix from 32 °C bath and allow to cool 1-2 degreesCritical: the mixture will start to solidify at 27-28 °C.
- 9 Add cell suspension to each tube, mix gently, then pour into plates.

- 10 Let solidify to a loose consistency
- 11 Transfer to growth room. Single colonies should appear within 10 days (without selection).