

Nov 12, 2023

Version 1

Preparation and Administration of Systemic 4-AP V.1

DOI

<https://dx.doi.org/10.17504/protocols.io.5qpvo38dxv4o/v1>

Leonardo Parra-Rivas¹

¹University of California, San Diego



Leonardo A Parra-Rivas

University of California, San Diego

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.5qpvo38dxv4o/v1>

Protocol Citation: Leonardo Parra-Rivas 2023. Preparation and Administration of Systemic 4-AP. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.5qpvo38dxv4o/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: November 12, 2023

Last Modified: November 12, 2023

Protocol Integer ID: 90816

Keywords: ap preparation, administration of systemic, ap, preparation, systemic

Abstract

Preparation and Administration of Systemic 4-AP

1

4-Aminopyridine was obtained from Sigma-Aldrich (Cat #A78403), and a stock solution was prepared in saline. Mice (8 weeks old) were given intraperitoneal injections of soluble 4-AP at a low dose of 2 mg/Kg or equivalent saline.

2 Animals were monitored every 20 min and presented no sign of convulsive seizure behavior.

After 3 hrs, animals were anesthetized with isoflurane (VetOne Cat#502017), followed by cervical dislocation. Brains were rapidly removed and drop-fixed in 4% PFA solution for immunohistochemistry or placed on ice for biochemical assays.