VERSION 1

JUN 14, 2017

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.iekcbcw

Collection Citation: Anibal Arce Medina, 2017.

Preparation of cell-free RNAPT7 reactions.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.iekcbcw>

MANUSCRIPT CITATION:

Anibal Arce, Juan Puig, Fernà Federici

<http://federicilab.org>

License: This is an open access collection distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Jun 12, 2017

Last Modified: Mar 17, 2018

COLLECTION integer ID:
6316

🌐 Preparation of cell-free RNAPT7 reactions V.1

Anibal Arce Medina,

Laboratorio de Tecnologías Libres

Anibal Arce Medina

Pontificia Universidad Catolica de Chile

ABSTRACT

This collection contains the protocols and recipes for the preparation of cell-free RNAPT7 transcription/translation coupled reactions.

The crude extracts are obtained from IPTG-treated E.coli BL21 DE3 STAR cells.

IPTG triggers RNAPT7 expression and accumulation in cells before collection. Reactions take place when the cell extract is mixed with a maltodextrin-based energy solution along with amino acids solution and other salts (Magnesium and Potassium). This system allows transcription/translation coupled reactions from plasmid DNA that contains T7 promoter transcriptional units. In our hands, it has been possible to express different fluorescent reporters, including deGFP which is highly efficient. Products of the reaction can be measured during the whole reaction time in a plate reader with appropriate optic filters, open hardware cameras, and microscopes. When set up correctly, deGFP fluorescence intensity is detected as fast as ~30 minutes after the reaction starts and it shows a rapid increase during the first ~2.5-3.0 hours.

The system presented here contains the features highlighted below, and is an adaptation of protocols and findings from the following listed papers:

1.- This system uses maltodextrin and Polyphosphate-based energy solution:

Kim, H.-C., Kim, T.-W., Kim, D.-M., . Prolonged production of proteins in a cellfree protein synthesis system using polymeric carbohydrates (2011)

Caschera, F.; Noireaux, V., A cost-effective polyphosphate-based metabolism fuels an all E. coli cell-free expression system. Metabolic engineering (2015)

2.- The S12 crude extract is prepared from E.coli cells (expressing Pol T7) using bead beater.

Sun, Z. Z.; Hayes, C. A.; Shin, J.; Caschera, F.; Murray, R. M.; Noireaux, V., Protocols for implementing an Escherichia coli based TX-TL cell-free expression system for synthetic biology. Journal of visualized experiments (2013)

Kim, TW., Kim, HC., Oh, IS. et al. A highly efficient and economical cell-free protein synthesis system using the S12 extract of Escherichia coli. Biotechnol Bioproc E (2008)

3.- The amino acid stock solution contains each amino acid at ~12 nM in a stable solution:

Caschera, V. Noireaux Preparation of amino acid mixtures for cell-free expression systems. Benchmarks (2015)

FILES

Protocol

NAME

Step 3: Performing cell-free RNAPT7 reactions

VERSION 1

CREATED BY

Anibal Arce Medina Pontificia Universidad Catolica de Chile

[OPEN](#) →

Protocol

NAME

Step 1: Preparing S12 cell extracts using bead-beater

VERSION 1

CREATED BY

Anibal Arce Medina Pontificia Universidad Catolica de Chile

[OPEN](#) →

Protocol

NAME

Step 2: Preparing amino acid, polyphosphates, and maltodextrin-based energy solutions for cell-free reactions

VERSION 1

CREATED BY

Anibal Arce Medina Pontificia Universidad Catolica de Chile

[OPEN](#) →