MAR 25, 2020

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.gtcbwiw

Protocol Citation: Alice Pawlowski 2020. Preparation of chemically competent *E. coli* cells. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.gtcbwiw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Dec 19, 2016

Last Modified: Mar 25, 2020

PROTOCOL integer ID:
 4676

Keywords: *E. coli*, transformation, competent cells, CaCl₂

Preparation of chemically competent *E. coli* cells

Alice Pawlowski¹

¹Institute of Synthetic Microbiology, Heinrich Heine University Düsseldorf

Axmann Lab

Alice Pawlowski

ABSTRACT

Standard protocol to prepare cells for highly efficient DNA uptake using CaCl₂ (Sambrook and Russell, Molecular Cloning: A Laboratory Manual, 2001).

MATERIALS

MATERIALS

LB-Medium (Lennox) vegetal Carl Roth Catalog
 #0155.1

- 10 mM CaCl₂
- sterile glycerole

- 1 Inoculate a 5 ml overnight-culture in LB-medium without antibiotics.
- 2 Inoculate 100 ml LB-medium with 1/100 volume of the overnight culture and incubate to an $OD_{600} = 0.5 - 0.6$ at 37 °C and 230 rpm.
- 3 Transfer cells to two 50 ml sterile falcon tubes and incubate on ice for 20 min.

From this step on, keep cells always cold.

00:20:00

- 4 Centrifuge for 5 min at 4500 rpm and 4 °C (pre-cool centrifuge) and discard the supernatant.

00:05:00

- 5 Carefully suspend each pellet in 20 ml of icecold 100 mM $CaCl_2$ and incubate on ice for 1 h.

01:00:00

- 6 Centrifuge for 5 min at 4500 rpm and 4 °C (pre-cool centrifuge) and discard the supernatant.

00:05:00

- 7 Carefully suspend each pellet in 2 ml icecold 85 mM $CaCl_2$ containing 15% Glycerol.

- 8 Pipette 100 to 200 μ l aliquots in 1.5 ml tubes (work on ice) and store at -80 °C. (you can shock-freeze the cells in liquid nitrogen or in dry ice prior to storage at -80°C, but it is not required)

For DNA transformation, thaw 100 μ l of the competent cells on ice.