SEP 22, 2021

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bqnumview

External link:
https://link.springer.com/protocol/10.1007/978-1-4939-9121-1_4

Protocol Citation: Anne Zemella, Theresa Richter, Lena Thoring, Stefan Kubick 2021. Preparation of Enhanced Orthogonal Aminoacyl-tRNA-Synthetase. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bqnumview>

MANUSCRIPT CITATION:
 Zemella A., Richter T., Thoring L., Kubick S. (2019) A Combined Cell-Free Protein Synthesis and Fluorescence-Based Approach to Investigate GPCR Binding Properties. In: Tiberi M. (eds) G Protein-Coupled Receptor Signaling. Methods in Molecular Biology, vol 1947. Humana Press, New York, NY.
https://doi.org/10.1007/978-1-4939-9121-1_4

Preparation of Enhanced Orthogonal Aminoacyl-tRNA-Synthetase

In 1 collection

Anne Zemella¹, Theresa Richter^{1,2}, Lena Thoring¹, Stefan Kubick¹

¹Cell-free and Cell-based Bioproduction, Fraunhofer Institute for Cell Therapy and Immunology (IZI), Branch Bioanalytics and Bioprocesses (IZI-BB), Potsdam, Germany;

²University of Potsdam, Potsdam, Germany

Springer Nature Books

Satyavati Kharde
 Springer Nature

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: Dec 12, 2020

Last Modified: Sep 22, 2021

PROTOCOL integer ID:
45492

Keywords: Cell-free protein synthesis, G protein-coupled receptor, Protein modification, Non-canonical amino acids, Amber suppression, Confocal laser scanning microscopy

ABSTRACT

This is part 3.1 of the "A Combined Cell-Free Protein Synthesis and Fluorescence-Based Approach to Investigate GPCR Binding Properties" collection of protocols: <https://www.protocols.io/view/a-combined-cell-free-protein-synthesis-and-fluores-bqntmven>

Collection Abstract: Fluorescent labeling of de novo synthesized proteins is in particular a valuable tool for functional and structural studies of membrane proteins. In this context, we present two methods for the site-specific fluorescent labeling of difficult-to-express membrane proteins in combination with cell-free protein synthesis. The cell-free protein synthesis system is based on Chinese Hamster Ovary Cells (CHO) since this system contains endogenous membrane structures derived from the endoplasmic reticulum. These so-called microsomes enable a direct integration of membrane proteins into a biological membrane. In this protocol the first part describes the fluorescent labeling by using a precharged tRNA, loaded with a fluorescent amino acid. The second part describes the preparation of a modified aminoacyl-tRNA-synthetase and a suppressor tRNA that are applied to the CHO cell-free system to enable the incorporation of a non-canonical amino acid. The reactive group of the non-canonical amino acid is further coupled to a fluorescent dye. Both methods utilize the amber stop codon suppression technology. The successful fluorescent labeling of the model G protein-coupled receptor adenosine A2A (Adora2a) is analyzed by in-gel-fluorescence, a reporter protein assay, and confocal laser scanning microscopy (CLSM). Moreover, a ligand-dependent conformational change of the fluorescently labeled Adora2a was analyzed by bioluminescence resonance energy transfer (BRET).

For **Introduction and Notes**, please see: <https://www.protocols.io/view/a-combined-cell-free-protein-synthesis-and-fluores-bqntmven/guidelines>

MATERIALS

2.1 Materials for Preparation of Enhanced Orthogonal Aminoacyl-tRNA-Synthetase

1. Coding sequence for the modified tyrosyl-tRNA-synthetase (eAzFRS, including the mutations Thr37, Ser182, Ala183, and Arg265 [11, 12] and a C-terminal Strep-Tag) from *E. coli*.
2. *E. coli* expression system (RTS 500 *E. coli* HY Kit, biotech rabbit).
3. 100 mM Isopropyl β -D-1-thiogalactopyranoside (IPTG)
4. Gravity flow Strep-Tactin[®] superflow mini-column (0.2 ml).
5. Strep-Tactin[®] Purification Buffer Set: 10 $\times$ Washing Buffer (1 M Tris-Cl, pH 8.0, 1.5 M NaCl, 10 mM EDTA), 10 $\times$ Elution Buffer (1 M Tris-Cl, pH 8.0, 1.5 M NaCl, 10 mM EDTA, 25 mM Desthiobiotin) and 10 $\times$ Regeneration Buffer (1 M Tris-Cl, 1.5 M NaCl, 10 mM EDTA, 10 mM HABA (hydroxyl-azophenyl-benzoic acid)).
6. Zeba[™] Spin Desalting Columns (7 K MWCO, 0.5 ml).
7. Amicon[®] Ultra Centrifugal Filters (10 K device, 0.5 ml).
8. Synthetase storage buffer: 50 mM HEPES pH 7.6, 10 mM KOAc, 1 mM MgCl₂, 4 mM DTT.
9. Thermomixer with a microtiter plate adapter and a RTS 500 adapter.

SAFETY WARNINGS

For hazard information and safety warnings, please refer to the SDS (Safety Data Sheet).

3.1 Preparation of Enhanced Orthogonal Aminoacyl-tRNA-S...

20h

- 1 For prokaryotic cell-free synthesis, the eAzFRS gene should be cloned into a vector containing a T7 promotor, ribosomal binding site, and T7 terminator such as pIX3.0, pIVEX2.3d, and pIVEX2.4d vectors or alternatively containing a T5 promotor such as pQE2 vectors as used in this protocol. eAzFRS is synthesized in a cell-free system using an *E. coli* lysate in a dialysis mode. A typical 1.1 ml reaction is composed of 0.525 mL *E. coli* lysate, 0.225 mL reaction mix, 0.27 mL amino acids without methionine, 30 μ L methionine, 11 μ L IPTG for the induction of the protein expression pQE2 vector, 39 μ L template containing 110 μ g plasmid DNA.
- 2 The surrounding feeding mixture contains 7990 μ L feeding mix, 110 μ L IPTG, 2650 μ L amino acids without methionine and 300 μ L methionine (see Note 2).

- 3 Fill the reaction solution into the reaction compartment (marked through the red lid).
- 4 Fill the feeding mix into the feeding chamber (marked through the colorless lid).
- 5 Insert the prepared chamber into the RTS 500 adapter in a thermomixer. The reaction time is 20:00:00 at 30 °C and a shaking speed of 900 rpm . 20h
- 6 For the separation of aggregated proteins from soluble eAzFRS a centrifugation step at 16000 x g, 4°C, 00:10:00 is recommended.
- 7 Equilibrate two Strep-Tactin columns with 400 µL 10× washing buffer and add 500 µL supernatant of the cell-free reaction to each column.
- 8 After the supernatant has completely entered the column, wash each column 5× with 200 µL washing buffer (see **Note 3**).
- 9 Elute the protein 6× with 100 µL elution buffer and collect the fractions.
- 10 Elution fractions containing the target protein are pooled.

- 11 Regenerate the column with 3× 1 mL 1× regeneration buffer and remove the regeneration buffer 2× with 800 µL 1× washing buffer. Store the column in 2 mL washing buffer at 4 °C.
- 12 The combined elution fractions are applied to Zeba™ Spin Desalting Columns to exchange the elution buffer of the strep-tag purification to a synthetase storage buffer. Therefore, remove the storage solution of the Zeba™ Spin Desalting Column by centrifugation at 1500 x g, 00:01:00. Add 300 µL synthetase storage buffer to the resin bed and centrifuge at 1500 x g, 00:01:00. Repeat this **step 2×**. Place the column in a new collection tube and apply 100 µL pooled synthetase solution to each column. Centrifuge at 2000 x g, 00:02:00 and collect the synthetase.
- 13 The concentration of the synthetase can be performed with Amicon® Ultra Centrifugal Filters. Add up to 500 µL synthetase solution to the concentrator and centrifuge at 14000 x g, 4°C, 00:10:00. Collect the concentrated sample and determine the concentration by NanoDrop measurement using the molecular mass (48.6 kDa) and the extinction coefficient (54.3) (*see Note 4*).
- 14 The synthetase can be stored at -80 °C after shock freezing in liquid nitrogen.