JUL 29, 2017

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.igacbse

External link:
<https://doi.org/10.1371/journal.pone.0181703>

Protocol Citation: ShiNan Dong 2017. Preparation of Genomic DNA of Microsporidia. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.igacbse>

MANUSCRIPT CITATION:
 Chen G, Wang W, Chen H, Dai W, Peng X, Li X, Tang X, Xu L, Shen Z (2017) Functional characterization of an aquaporin from a microsporidium, *Nosema bombycis*. PLoS ONE 12(7): e0181703. doi: [10.1371/journal.pone.0181703](https://doi.org/10.1371/journal.pone.0181703)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Preparation of Genomic DNA of Microsporidia

PLOS One

ShiNan Dong

Chen Gong

ABSTRACT

To prepare the genomic DNA of microsporidia, 400 μ l suspension of purified spores (1010 spores/ml) was mixed with 40 μ l KOH (2 mol/l) in a 1.5 ml Eppendorf tube and incubated at 27°C for 1 h, TEK buffer (1 mmol/l Tris-HCl, 10 mmol/l EDTA, 0.17 mol/l KCl, pH 8.0) was added and continued incubating at 27°C for 1 h. Adjusted pH to 8.0 with 1 mol/l HCl, added 10% SDS in order to attain 0.5% in the mixture and kept it in ice-bath for 15 min. 20 mg/ml proteinase K (TaKaRa Biotechnology Co. LTD) was added in order to attain 200 μ g/ml in the mixture and incubated at 50°C for 4 h. Subsequently, isovolumetric tris-phenol extraction was performed twice, and washed with chloroform: isoamyl alcohol (24:1 v/v). 10% NaOAC was added to the recovered aqueous phase after centrifuging (10,000 r/min, 5 min), and then the DNA was precipitated by 2.5 times volume of cold ethanol at -20°C for 40 min before centrifuging (12,000 r/min, 10 min). In addition, the precipitated DNA was rinsed twice with 500 μ l cold 70% ethanol before drying at 37°C for 5–10 min. Then, the extracted DNA was stored at -20°C after dissolution in 50 μ l TE buffer at 65°C for 10 min.

ATTACHMENTS

[Sequence and Phylogenetic Analysis of SSU rRNA Gene.pdf](#)

Created: Jun 14, 2017

Last Modified: Mar 14, 2018

PROTOCOL integer ID:
6370