AUG 22, 2020

🌐 Preparation of staphylococcal protein-A conjugated to horseradish peroxidase.

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bj5ykq7w

Protocol Citation: Angel A Justiz-Vaillant 2020. Preparation of staphylococcal protein-A conjugated to horseradish peroxidase.. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bj5ykq7w>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Aug 22, 2020

Last Modified: Aug 22, 2020

PROTOCOL integer ID:
 40856

ABSTRACT

This reagent can be used in ELISA, Western blotting and Dot blot to detect antigens and antibodies. It is important in the immunodiagnosis of infectious diseases. I find this useful in the detection of anti-HIV antibodies by ELISA.

MATERIALS

MATERIALS

⊗ Ammonium Sulfate **P212121**

⊗ Sodium periodate **Bio Basic Inc. Catalog #SB0875.SIZE.100g**

⊗ sodium borohydride **Sigma Aldrich Catalog #452882**

⊗ Horseradish Peroxidase (HRP) type IV **Sigma Aldrich Catalog #P8375-25KU**

⊗ Staphylococcal Protein-A **Sigma Aldrich**

Pipettes

20ml to 1000 ml glass

Scale

Incubator

Refrigerator

Freezer

Centrifuges

- 1 Horseradish peroxidase (500 µg in 50 µl NaCO₃, pH 9.6) is mixed with freshly made sodium periodate solution (1.71 mg/ml) followed by incubation in the dark for 2 h.

- 2 Mix 500 µg of staphylococcal protein-A (SpA) with an equal amount (500 micrograms) of a mix of horseradish peroxidase-sodium periodate.
- 3 The mixture is incubated for 3 hours at 4°C with gentle agitation.
- 4 Forty µl of freshly prepared NaBH₄ solution (5 mg NaBH₄ /ml 0.1 mM NaOH) is then added to the preparation.
- 5 The preparation is incubated for 90 min at 4°C in the dark with gentle agitation.
- 6 Cold 50% saturated ammonium sulphate solution (pH 7.4) is added drop by drop in the ratio 1:1 (v/v).
- 7 The mixture is then centrifuged for 25 min at 4°C and recover the pellet at the bottom of the tube.
- 8 The pellets is re-suspended in 200 µl of PBS pH=7.4 and dialysed against 1L of PBS for 24 h with 3 buffer changes.
- 9 An equal volume of glycerol is added to the dialysate followed by 100 µl of bovine serum albumin, BSA (20 mg/ ml).

10 The conjugate is then stored at -20°C until further used.