

Aug 12, 2020

Preparation of staphylococcal protein-A conjugated to horseradish peroxidase by the periodate method.

DOI

dx.doi.org/10.17504/protocols.io.bjk5kky6

Angel A Justiz-Vaillant¹

¹University of the West Indies St. Augustine

University of the West In...

angel.vaillant@sta.uwi.e...



Angel A Justiz-Vaillant

University of the West Indies St. Augustine

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bjk5kky6

Protocol Citation: Angel A Justiz-Vaillant 2020. Preparation of staphylococcal protein-A conjugated to horseradish peroxidase by the periodate method.. [protocols.io https://dx.doi.org/10.17504/protocols.io.bjk5kky6](https://dx.doi.org/10.17504/protocols.io.bjk5kky6)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: August 12, 2020

Last Modified: August 12, 2020

Protocol Integer ID: 40317

Abstract

This reagent can be used in ELISA, Western blotting and Dot blot to detect antigens and antibodies. It is important in the immunodiagnosis of infectious diseases and other problems. I find this useful in the detection of anti-HIV antibodies by ELISA.

Guidelines

All reagents but specially the enzyme and sodium periodate solution has to be prepared freshly before mixing it with the enzyme.

Materials

MATERIALS

- ⊗ Ammonium Sulfate **P212121**
- ⊗ Sodium periodate **Bio Basic Inc. Catalog #SB0875.SIZE.100g**
- ⊗ sodium borohydride **Sigma Aldrich Catalog #452882**
- ⊗ Horseradish Peroxidase (HRP) type IV **Sigma Aldrich Catalog #P8375-25KU**
- ⊗ Staphylococcal Protein-A **Sigma Aldrich**

Pipettes

20ml to 1000 ml glass

Scale

Incubator

Refrigerator

Freezer

Centrifuges

- 1 Horseradish peroxidase (500 µg in 50 µl NaCO₃ , pH 9.6) is mixed with freshly made sodium periodate solution (1.71 mg/ml) followed by incubation in the dark for 2 h.
- 2 Mix 500 µg of staphylococcal protein-A (SpA) with an equal amount (500 micrograms) of a mix of horseradish peroxidase-sodium periodate.
- 3 The mixture is incubated for 3 hours at 4°C with gentle agitation.
- 4 Forty µl of freshly prepared NaBH₄ solution (5 mg NaBH₄ /ml 0.1 mM NaOH) is then added to the preparation.
- 5 The preparation is incubated for 90 min at 4°C in the dark with gentle agitation.
- 6 Cold 50% saturated ammonium sulphate solution (pH 7.4) is added drop by drop in the ratio 1:1 (v/v).
- 7 The mixture is then centrifuged for 25 min at 4°C and recover the pellet at the bottom of the tube.
- 8 The pellets is re-suspended in 200 µl of PBS pH=7.4 and dialysed against 1L of PBS for 24 h with 3 buffer changes.
- 9 An equal volume of glycerol is added to the dialysate followed by 100 µl of bovine serum albumin, BSA (20 mg/ ml).
- 10 The conjugate is then stored at -20°C until further used.

11