VERSION 1

JUN 03, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3nrgmd6

Protocol Citation: Cancer Research UK / Wellcome Gurdon Institute media kitchen 2019. Prepare NGM no peptone plates. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3nrgmd6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 03, 2019

Last Modified: Jun 03, 2019

PROTOCOL integer ID:
 23985

1

NGM No Peptone agar_5L.xls

🌐 Prepare NGM no peptone plates V.1

Cancer Research UK / Wellcome Gurdon Institute media kitchen¹

¹Cancer Research UK / Wellcome Gurdon Institute

Cristian Riccio
 University of Cambridge

ABSTRACT

Prepare NGM plates without peptone.

Peptone is necessary for bacterial growth and thus by not adding peptone, one can control for the number of bacteria present on an NGM plate by seeding a known number of bacteria.

MATERIALS

MATERIALS

Agar **Contributed by users**

double distilled water (ddH2O) **Contributed by users**

Sodium chloride meets analytical specification of Ph.Eur Fisher Chemical **Fisher Scientific Catalog #S/3160/65**

SYCHEM autoclave **Syschem**

SAFETY WARNINGS

Make sure you know how to use the autoclave before starting this protocol.

2

Ingredients	Quantity	
NaCl		15g
Double distilled H2O	4.8L	
Agar	per 1L bottle	17g
Agar	per 500ml bottle	8.5g

3

Measure approx 4.8L double distilled H2O in 5L bell jar with a magnetic flea.
Add NaCl
Stir until all solutes are dissolved.
Dispense approx 972ml NGM media per 1L bottle OR 486ml NGM into a 500ml bottle.
Dispense 17g of agar and a magnetic flea to each 1L bottle OR 8.5g of agar and a magnetic flea to each 500ml bottle.
Label, date and autoclave.