VERSION 1

JUN 03, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.3ntgmen

Protocol Citation: Cancer Research UK / Wellcome Gurdon Institute media kitchen 2019. Prepare NGM plates for nematode, with peptone, without fungizone.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.3ntgmen>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jun 03, 2019

Last Modified: Jun 03, 2019

PROTOCOL integer ID:
 23987

🌐 Prepare NGM plates for nematode, with peptone, without fungizone V.1

Cancer Research UK / Wellcome Gurdon Institute media kitchen¹

¹Cancer Research UK / Wellcome Gurdon Institute

Cristian Riccio
 University of Cambridge

ABSTRACT

Prepare NGM plates for nematodes, with peptone, without fungizone

MATERIALS

MATERIALS

⊗ Agar **Contributed by users**

⊗ Difco Bacto peptone **Fisher Scientific**

⊗ double distilled water (ddH2O) **Contributed by users**

⊗ Sodium chloride meets analytical specification of Ph.Eur Fisher Chemical **Fisher Scientific Catalog #S/3160/65**

⊗ SYCHEM autoclave **Syschem**

SAFETY WARNINGS

Make sure you know how to use the autoclave before starting this protocol.

1

NGM_agar_1L.xls

2

Ingredients		Quantity
NaCl		3g
bacto peptone		2.5g
Agar		17g
Double distilled H2O	972ml	

3

Measure 972ml double distilled H2O and put in a 1L duran bottle with a magnetic flea
Add 3g NaCl, 2.5g bacto peptone and stir.
Stir until all solutes are dissolved then add the agar.
Leave magnetic flea in bottle.
Label, date and autoclave