VERSION 1

SEP 14, 2019

Prepare Uric Acid Solution V.1

宏亮 董¹

¹Northeast Forest University

2019 iGEM NEFU_China

Tech. support email: shengyiyanwork@gmail.com

宏亮 董

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.7cjhiun

Protocol Citation: 宏亮 董

2019. Prepare Uric Acid Solution. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.7cjhiun>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In

development

We are still developing and optimizing this protocol

Created: Sep 14, 2019

Last Modified: Sep 14, 2019

PROTOCOL integer ID:

27755

Keywords: Uric acid, storage solution

Prepare **Uric Acid Storage Solution:**

1 Accurately weigh 0.01g uric acid dissolved in **Uric Acid Diluent Solution** and dilute to 100mL;

2 Prepare **Borate Buffer Storage Solution**:

Add these reagents to a clean media bottle, DDW volume to 200ml, adjust pH to 8.5 with NaOH.

boric acid	6.185 g
borax	9.535 g
EDTA	1.169 g
10% Triton X-100	400µl

3 Prepare **Uric Acid Diluent Solution**:

Add 152 mL DDW and 8 mL **Borate Buffer Storage Solution** to a clean media bottle, adjust pH to 8.5 with NaOH.

4 Prepare **Uric Acid Working Solution**:

Dilute **Uric Acid Storage Solution** with **Uric Acid Diluent Solution** for 10 times.

The concentration can be adjusted according to experimental requirements.