MAR 20, 2020

Preparing 1/2 SŠ Algal Inorganic Nutrient Medium

In 1 collection

Jakub Nedbal¹, Lu Gao^{2,3}

¹King's College London; ²Forschungszentrum Jülich;

³Heinrich Heine University

Jakub Nedbal

King's College London

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.bdza72e

External link:
<https://app.labstep.com/sharelink/ddc19dd9-4958-42fb-818d-9238c4384ec8>

Protocol Citation: Jakub Nedbal, Lu Gao 2020. Preparing 1/2 SŠ Algal Inorganic Nutrient Medium. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bdza72e>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development
We are still developing and optimizing this protocol

Created: Mar 20, 2020

Last Modified: May 29, 2020

PROTOCOL integer ID:
34562

ABSTRACT

This protocol describes preparation of 1/2 SŠ inorganic nutrient medium for *C. vulgaris*, *D. quadricauda*, and other fresh water microalgae. The medium is prepared out of sterile components inside sterile laminar flow hood. The medium recipe is based on [\[Zachleder & Šetlík, 1982\]](#).

GUIDELINES

This protocol describes preparation of a culture medium derived from [1/2 SŠ inorganic nutrient medium](#) [\[Zachleder & Šetlík, 1982\]](#) for growth of fresh water microalgae on a shaking orbital incubator. The solution of the 1/2 SŠ medium (0.5 g/l) is prepared by mixing stocks of solutions of individual components in autoclaved deionized water. The final solution pH is adjusted by KOH and filter sterilized.

The protocol described here differs slightly from the source recipe. The original medium recipe is aimed at producing up to 6 g/l (dry weight) culture density. We use the cells for imaging and there is no need to produce such dense cultures. Therefore the nutrients were diluted 12 times in this protocol and the medium can only support cultures up to 0.5 g/l (dry weight) culture density. The original recipe does not include source of carbon, assuming sufficient gas exchange and reliance on atmospheric CO₂. The flask-based culture conditions limit atmospheric gas exchange. This protocol therefore adds 0.83 mM NaHCO₃ as a source of carbon.

The whole process should take about 1 hour.

Keywords: Growth medium, inorganic growth medium, culture medium, microalgae, *C. vulgaris*, *D. quadricauda*

MATERIALS

Autoclaved deionized water in glass bottle (0.5 l)

1/2 SŠ Stock KNO_3 (1.83 M, filter sterilized)

1/2 SŠ Stock KH_2PO_4 (174 mM, filter sterilized)

1/2 SŠ Stock $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (82.8 mM, filter sterilized)

Sodium Bicarbonate (500 mM, filter sterilized)

1/2 SŠ Stock $\text{Fe} \cdot \text{Na} \cdot \text{EDTA}$ (9.1 μM , filter sterilized)

1/2 SŠ Stock CaCl_2 (66 μM , filter sterilized)

1/2 SŠ Stock Trace Elements (83 $\mu\text{l/l}$, filter sterilized)

[HI-98100 Checker Plus pH Tester](#) or other pH meter. Make sure the pH meter is calibrated before its use.

Potassium Hydroxide (2M, filter sterilized) for pH titration.

15 ml Falcon tube

Pipettes and tips

SAFETY WARNINGS

The constituents of the medium are chemicals, which must be handled and stored according to local regulations. Potassium hydroxide (KOH) is caustic and stands out in its health and safety risks. Handle KOH with care. Avoid spilling it. Protect your eyes and hands with suitable personal protective equipment. Wipe away any spillage.

BEFORE START INSTRUCTIONS

Ensure that all components are prepared and filter-sterilized. Autoclave deionized water in a glass bottle.

This protocol adds sodium bicarbonate, which is not in the [original protocol](#) and which was added as source of carbon for the algae. Autoclaving destroys sodium bicarbonate. It causes a chemical reaction converting the bicarbonate to into carbonate and change in pH.

Preparatory Steps

- 1 Prepare the components listed in the following table. Make sure all constituents have been sterilized and that the deionized water has been autoclaved.

Work in a sterile laminar flow hood.

Component	Quantity	Unit	Note
Final Volume	503	ml	
Maximum Dry Culture Weight	0.5	g/l	
Deionized Water	500	ml	Autoclaved
1/2 SŠ Stock KNO ₃ (1.83 M)	833	μl	
1/2 SŠ Stock KH ₂ PO ₄ (184 mM)	417	μl	
1/2 SŠ Stock MgSO ₄ ·7H ₂ O (82.8 mM)	417	μl	
Sodium Bicarbonate (500 mM)	833	μl	Do not autoclave
1/2 SŠ Stock Fe·Na·EDTA (9.1 μM)	417	μl	Do not autoclave
1/2 SŠ Stock CaCl ₂ (66 μM)	417	μl	
1/2 SŠ Trace Elements (1 ml/l)	42	μl	
2M KOH	20	μl	The goal is to get pH 7.5

Constituents of the 1/2 SŠ inorganic nutrient medium.

- 2 Autoclave 500 ml of deionized water in a glass clean bottle.

Let it cool down.

- 3 Prepare all the reagents and labware.

Medium Preparation

- 4 Work in a sterile laminar flow hood.

Add components listed in table in STEP 1 into the autoclaved deionized water in the order listed in the table. Do not add KOH yet.

- 5 Adjust the pH to 7.5 using KOH.

- 5.1 Take a small aliquot (1.5 ml) of the medium into a 15 ml Falcon tube.

- 5.2 Measure the pH of the medium inside the Falcon tube using the pH meter.

- 5.3 Empty the Falcon tube.

- 5.4 Add a small aliquot (5 μ l) of 2 M KOH to the medium. Mix well by closing the bottle and shaking or turning up side down a few times.

- 5.5 Repeat by going back to STEP 5.1, until pH reaches a value of approximately 7.5.

6 Label the bottle (1/2 SŠ, initials, date, pH).

7 Store at 4 °C in dark.

Warm up to room temperature each time before use.

Only open in a sterile cabinet.

Summary

8 This protocol describes the preparation of a medium for growth of freshwater algae on an orbital shaking incubator. The medium is to be used to dilute the growing cultures at regular intervals. The medium needs to be stored in a fridge at 4 °C and warmed up each time to room temperature to minimize the stress to the culture.

References

Protocol for preparing the medium using filter sterilization: [Preparing 1/2 SŠ algal inorganic nutrient medium](#)

[1/2 SŠ inorganic nutrient medium](#): Recipe from the [Culture Collection of Autotrophic Organisms](#)

[Zachleder & Šetlík, 1982](#): The original article introducing this medium