VERSION 2

JUL 26, 2019

OPEN ACCESS

DOI:
dx.doi.org/10.17504/protocols.io.5wig7ce

Protocol Citation: Gurdon Institute media kitchen 2019. Preparing 10 L of M9 buffer for nematode culture. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.5wig7ce>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
 We use this protocol and it's working

Created: Jul 26, 2019

Last Modified: Jul 26, 2019

PROTOCOL integer ID:
 26282

Keywords: M9, m9, nematodes, worms

🌐 Preparing 10 L of M9 buffer for nematode culture V.2

Gurdon Institute media kitchen¹

¹Cancer Research UK / Wellcome Gurdon Institute

Cristian Riccio
 University of Cambridge

ABSTRACT

Prepare 10 litres of M9 solution for collecting and washing *C. elegans*.

MATERIALS

MATERIALS

⊗ double distilled water (ddH2O) **Contributed by users**

⊗ Di-Sodium Hydrogen Orthophosphate Dihydrate Certified AR for Analysis Fisher Chemical **Contributed by users Catalog #S/4450/53**

⊗ Sodium chloride meets analytical specification of Ph.Eur Fisher Chemical **Fisher Scientific Catalog #S/3160/65**

⊗ Magnesium Sulfate Heptahydrate Certified AR for Analysis Fisher Chemical **Fisher Scientific Catalog #M/1050/53**

⊗ Potassium Dihydrogen Orthophosphate Certified AR for Analysis Fisher Chemical **Fisher Scientific Catalog # P/4800/53**

⊗ SYCHEM autoclave **Syschem**

SAFETY WARNINGS

⚠ This protocol makes use of an autoclave. Make sure you know how to use it properly before starting the protocol.

1

20m

Ingredients	Quantity	
Na ₂ HPO ₄		60g
KH ₂ PO ₄		30g
NaCl		50g
Double distilled H ₂ O	up to 10L	
Add 1M MgSO ₄ after autoclaving @ 100ul/100ml

M9_buffer_10L.xls

M9_buffer_2L.xls

2

15m

1	Dissolve ingredients in 8L double distilled H ₂ O
2	Adjust volume to 10L using double distilled H ₂ O
3	Measure or use pump to dispense aliquots accurately.
4	Label, date and autoclave.
5	After autoclaving, add 100ul 1M MgSO ₄ per 100ml. (ie. 100ul to 100ml and 200ul to 200ml bottles).
NB	1M MgSO ₄ at 1ul/1ml in the M9 buffer gives a final concentration of 1mM

3 The autoclave is SYCHEM VS Series. The media cycle is 15 minutes at 121 degrees centigrade. It is then cooled down in the machine for 3 hours. 3h 15m

4 After autoclaving, add 100 µl 1M MgSO₄ per 100 ml. (ie. 100 µl to 100 ml and 200 µl to 200 ml bottles). 10m

NB: 1M MgSO₄ at 1 µl/1ml in the M9 buffer gives a final concentration of 1 mM