



Sep 06, 2022

Primary neuronal cultures

 [Nature Communications](#)

DOI

<https://dx.doi.org/10.17504/protocols.io.ewov1ojwklr2/v1>

Miguel Da Silva Padilha¹, Irina Dudanova¹, F. Ulrich Hartl¹, Itika Saha², Mark S. Hipp^{3,4}

¹Molecular Neurodegeneration Group, Max Planck Institute of Neurobiology, 82152 Martinsried, Germany;

²Department of Cellular Biochemistry, Max Planck Institute of Biochemistry, Am Klopferspitz 18, 82152 Martinsried, Germany;

³Department of Biomedical Sciences of Cells and Systems, University Medical Center Groningen, University of Groningen, Antonius Deusinglaan, 1, 9713 AV Groningen, The Netherlands;

⁴School of Medicine and Health Sciences, Carl von Ossietzky University Oldenburg, 26129 Oldenburg, Germany



Felix Kraus

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.ewov1ojwklr2/v1>

External link: <https://www.biorxiv.org/content/10.1101/2022.02.18.481043v1.full>



Protocol Citation: Miguel Da Silva Padilha, Irina Dudanova, F. Ulrich Hartl, Itika Saha, Mark S. Hipp 2022. Primary neuronal cultures. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.ewov1ojwklr2/v1>

Manuscript citation:

Itika Saha, Patricia Yuste-Checa, Miguel Da Silva Padilha, Qiang Guo, Roman Körner, Hauke Holthusen, Victoria A. Trinkaus, Irina Dudanova, Rubén Fernández-Busnadiego, Wolfgang Baumeister, David W. Sanders, Saurabh Gautam, Marc I. Diamond, F. Ulrich Hartl, Mark S. Hipp bioRxiv 2022.02.18.481043; doi: <https://doi.org/10.1101/2022.02.18.481043>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: September 05, 2022

Last Modified: May 31, 2024

Protocol Integer ID: 69593

Keywords: ASAPCRN, preparation of primary neuronal culture, primary neuronal cultures this protocol, primary neuronal culture, neuronal culture, wild type mouse embryo, culture

Abstract

This protocol describes the preparation of primary neuronal cultures from E15.5 CD-1 wild type mouse embryos. Experiments involving animal models must be performed in accordance with relevant institutional guidelines and regulations.



- 1 Sacrifice pregnant female mice by cervical dislocation.
- 2 Remove the uterus from the abdominal cavity and place into a 10 cm sterile Petri dish on ice containing dissection medium, consisting of Hanks' balanced salt solution (HBSS) supplemented with 0.01 M HEPES, 0.01 M MgSO₄ and 1% penicillin/streptomycin.
- 3 Isolate each embryo, decapitate the heads, remove the brains from the skull and immerse in ice-cold dissection medium.
- 4 Dissect cortical hemispheres, and remove meninges under a dissection microscope.
- 5 Collect the cortices in a 15 mL sterile tube and digest with 0.25% trypsin containing 1 mM ethylenediaminetetraacetic acid (EDTA) and 15 μ L 0.1% DNase I for 20 min at 37 °C.
- 6 Stop digestion by removing the supernatant and washing the tissue twice with Neurobasal medium (Invitrogen) containing 5% Fetal Bovine Serum.
- 7 Resuspend the tissue in 2 mL Neurobasal medium and triturate to achieve a single cell suspension.
- 8 Spin cells at 130 x g, remove the supernatant, and resuspend the cell pellet in Neurobasal medium with 2% B-27 supplement (Invitrogen), 1% L-glutamine (Invitrogen) and 1% penicillin/streptomycin (Invitrogen).
- 9 Plate cells at desired density in dishes or coverslips coated with 1 mg/mL poly-D-lysine (Sigma) and 1 μ g/mL laminin (Thermo Fisher Scientific).