

Jan 02, 2024

 Prognostic value of combining 24-hour ASPECTS, and hemoglobin to red cell distribution width ratio on the THRIVE-score in predicting in-hospital mortality among ischemic stroke patients treated with intravenous thrombolysis.

DOI

<https://dx.doi.org/10.17504/protocols.io.dm6gp3o5pvzp/v1>

Sarawut Krongsut¹, Wipasiri Naraphong², Surachet Srikaew³, Niyada Anusasnee⁴

¹Division of Neurology, Department of Internal Medicine, Faculty of Medicine Saraburi;

²Department of Adult and Gerontological Nursing, Boromarajonani College of Nursing, Saraburi, Faculty of Nursing, Praboromarajchanok Institute, Ministry of Public Health, Saraburi, Thailand;

³Division of Neurosurgery, Department of Surgery, Faculty of Medicine, Srinakharinwirot University, Ongkharak Campus, Nakhon Nayok, Thailand;

⁴Division of radiology, Saraburi Hospital, Saraburi, Thailand



สราวุธ กรองสัตย์

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.dm6gp3o5pvzp/v1>

Protocol Citation: Sarawut Kongsut, Wipasiri Naraphong, Surachet Srikaew, Niyada Anusasnee 2024. Prognostic value of combining 24-hour ASPECTS, and hemoglobin to red cell distribution width ratio on the THRIVE-score in predicting in-hospital mortality among ischemic stroke patients treated with intravenous thrombolysis.. protocols.io

<https://dx.doi.org/10.17504/protocols.io.dm6gp3o5pvzp/v1>

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 19, 2023

Last Modified: January 02, 2024

Protocol Integer ID: 89613

Keywords: ASPECTS, thrombolysis, in-hospital mortality, ischemic stroke, predictive value, precise prediction of stroke outcome, stroke outcome, stroke clinical outcome, hospital mortality among ischemic stroke patient, acute ischemic stroke, ischemic stroke patient, hemoglobin to red cell distribution width ratio, total health risk in vascular event, hour alberta stroke program, hours of intravenous thrombolysis completion, ratio into the thrive score, intravenous thrombolysis completion, prognostic value, stroke, data on thrive score, thrive score, clinical outcome, receiving intravenous thrombolysi, intravenous thrombolysi, middle cerebral artery territory, ischemic cerebral lesion, total health risk, hemoglobin, hospital mortality, impacting mortality, red cell distribution width ratio

Funders Acknowledgements:

Medical Education Center at Saraburi Hospital

Grant ID: MC013-2566

Disclaimer

The authors have no conflicts of interest directly relevant to the content of this article.

Abstract

Background:

Acute ischemic stroke (AIS) emerges as a significant global health issue, directly impacting mortality and disability. The Total Health Risk In Vascular Events (THRIVE) score, appreciated for its simplicity and ease of use, consistently validates its ability to accurately predict stroke clinical outcomes. However, the THRIVE score is currently lacking laboratory and neuroimaging data, which limits its ability to provide precise prediction of stroke outcomes. Our study evaluates the impact of integrating the 24-hour Alberta Stroke Program Early CT Score (ASPECTS) and hemoglobin-to-red cell distribution width (HB/RDW) ratio into the THRIVE score using the multivariable fractional polynomial (MFP) method (combined THRIVE-MFP) compared to the THRIVE-c model. We aim to assess their added value in predicting in-hospital mortality (IHM) prognosis.

Methods: A retrospective study from January 2015 to July 2022 examined consecutive AIS patients receiving intravenous thrombolysis. Data on THRIVE scores, 24-hour ASPECTS, and HB/RDW levels were collected upon admission. The model was constructed using logistic regression and the MFP method. The prognostic value was determined using the augmented area under the receiver operating characteristic (AuROC). Ischemic cerebral lesions within the middle cerebral artery territory were evaluated with non-contrast computed tomography (NCCT) following 24 hours of intravenous thrombolysis completion (24-hour ASPECTS).

Results:

Among a cohort of 345 patients diagnosed with AIS who received intravenous thrombolysis, 65 individuals (18.8%) experienced IHM. The combined THRIVE-MFP model was significantly superior to the THRIVE-c model (AuROC 0.980 vs. 0.876, $P < 0.001$).

Conclusion:

The combined THRIVE-MFP model showed excellent predictive performance, offering potential for guiding treatment decisions and enhancing healthcare outcomes. Incorporating 24-hour ASPECTS on NCCT and HB/RDW proved valuable in IHM prediction, particularly for hospitals with limited access to advanced neuroimaging resources.

Attachments



[Protocol.pdf](#)

211KB



- 1 Prognostic value of combining 24-hour ASPECTS, and hemoglobin to red cell distribution width ratio on the THRIVE-score in predicting in-hospital mortality among ischemic stroke patients treated with intravenous thrombolysis

Protocol references

1. Langhorne P, Dey P, Woodman M, et al. Is stroke unit care portable? A systematic review of the clinical trials. *Age Ageing*. 2005;34(4):324-330. doi:10.1093/ageing/afi038
2. Krishnamurthi R V., Ikeda T, Feigin VL. Global, Regional and Country-Specific Burden of Ischaemic Stroke, Intracerebral Haemorrhage and Subarachnoid Haemorrhage: A Systematic Analysis of the Global Burden of Disease Study 2017. *Neuroepidemiology*. 2020;54(2):171-179. doi:10.1159/000506396
3. Phongphuttha W, Tiamkao S. Outcome of stroke fast-track patients arrival by emergency medical services. *J Med Assoc Thail*. 2021;104(2):S88-S93. doi:10.35755/jmedassothai.2021.S01.12299
4. Chatterjee S. Recombinant tissue plasminogen activator for acute ischemic stroke. *Cardiol Rev*. 2012;28(4).
5. Counsell C, Dennis M, McDowall M, Warlow C. Predicting outcome after acute and subacute stroke: Development and validation of new prognostic models. *Stroke*. 2002;33(4):1041-1047. doi:10.1161/hs0402.105909
6. Johnston KC, Wagner DP, Wang XQ, et al. Validation of an acute ischemic stroke model: Does diffusion-weighted imaging lesion volume offer a clinically significant improvement in prediction of outcome? *Stroke*. 2007;38(6):1820-1825. doi:10.1161/STROKEAHA.106.479154
7. Wang Y, Lim LLY, Levi C, F Heller R, Fischer J. A prognostic index for 30-day mortality after stroke. *J Clin Epidemiol*. 2001;54(8):766-773. doi:10.1016/S0895-4356(01)00338-9
8. Flint AC, Cullen SP, Faigles BS, Rao VA. Predicting long-term outcome after endovascular stroke treatment: The totaled health risks in vascular events score. *Am J Neuroradiol*. 2010;31(7):1192-1196. doi:10.3174/ajnr.A2050
9. Ader F, Bouscambert-Duchamp M, Hites M, et al. Remdesivir plus standard of care versus standard of care alone for the treatment of patients admitted to hospital with COVID-19 (DisCoVeRy): a phase 3, randomised, controlled, open-label trial. *Lancet Infect Dis*. 2022;22(2):209-221. doi:10.1016/S1473-3099(21)00485-0
10. Flint AC, Rao VA, Chan SL, et al. Improved ischemic stroke outcome prediction using model estimation of outcome probability: The THRIVE-c calculation. *Int J Stroke*. 2015;10(6):815-821. doi:10.1111/ijss.12529
11. Barber PA, Demchuk AM, Zhang J, Buchan AM. Validity and reliability of a quantitative computed tomography score in predicting outcome of hyperacute stroke before thrombolytic therapy. *Lancet*. 2000;355(9216):1670-1674. doi:10.1016/S0140-6736(00)02237-6
12. Kong WY, Tan BYQ, Ngiam NJH, et al. Validation of Serial Alberta Stroke Program Early CT Score as an Outcome Predictor in Thrombolized Stroke Patients. *J Stroke Cerebrovasc Dis*. 2017;26(10):2264-2271. doi:10.1016/j.jstrokecerebrovasdis.2017.05.009
13. Liebeskind DS, Jahan R, Nogueira RG, Jovin TG, Lutsep HL, Saver JL. Serial Alberta stroke program early CT score from baseline to 24 Hours in solitaire flow restoration with the intention for thrombectomy study: A Novel surrogate end point for revascularization in acute stroke. *Stroke*. 2014;45(3):723-727. doi:10.1161/STROKEAHA.113.003914
14. Mitomi M, Kimura K, Aoki J, Iguchi Y. Comparison of CT and DWI findings in ischemic stroke patients within 3 hours of onset. *J Stroke Cerebrovasc Dis*. 2014;23(1):37-42. doi:10.1016/j.jstrokecerebrovasdis.2012.08.014
15. Turcato G, Cervellin G, Cappellari M, et al. Early function decline after ischemic stroke can be predicted by a nomogram based on age, use of thrombolysis, RDW and NIHSS score at admission. *J Thromb Thrombolysis*. 2017;43(3):394-400. doi:10.1007/s11239-016-1456-y

16. Mohindra R, Mishra U, Mathew R, Negi NS. Red Cell Distribution Width (RDW) Index as a Predictor of Severity of Acute Ischemic Stroke: A Correlation Study. *Adv J Emerg Med.* 2020;4(2):e24. doi:10.22114/ajem.v0i0.257
17. Rahamim E, Zwas DR, Keren A, et al. The Ratio of Hemoglobin to Red Cell Distribution Width: A Strong Predictor of Clinical Outcome in Patients with Heart Failure. *J Clin Med.* 2022;11(3). doi:10.3390/jcm11030886
18. Sun P, Zhang F, Chen C, et al. The ratio of hemoglobin to red cell distribution width as a novel prognostic parameter in esophageal squamous cell carcinoma: A retrospective study from southern China. *Oncotarget.* 2016;7(27):42650-42660. doi:10.18632/oncotarget.9516
19. Wang J, Chen Z, Yang H, Li H, Chen R, Yu J. Relationship between the Hemoglobin-to-Red Cell Distribution Width Ratio and All-Cause Mortality in Septic Patients with Atrial Fibrillation: Based on Propensity Score Matching Method. *J Cardiovasc Dev Dis.* 2022;9(11):341-354. doi:10.3390/jcdd9110400
20. Powers WJ, Rabinstein AA, Ackerson T, et al. Guidelines for the Early Management of Patients with Acute Ischemic Stroke: 2019 Update to the 2018 Guidelines for the Early Management of Acute Ischemic Stroke a Guideline for Healthcare Professionals from the American Heart Association/American Stroke A. Vol 50.; 2019. doi:10.1161/STR.0000000000000211