

Oct 06, 2019

Propagation and attenuation of FAdV isolate in CEL cells

 [PLOS One](#)

DOI

<https://dx.doi.org/10.17504/protocols.io.7zdhp26>

Norfitriah Mohamed Sohaimi¹, Mohd Hair Bejo¹, Abdul Rahman Omar¹, Aini Ideris¹, Nurulfiza Mat Isa²

¹Faculty of Veterinary Medicine, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.;

²Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, Serdang, Selangor, Malaysia.



Norfitriah Mohamed Sohaimi

Create & collaborate more with a free account

Edit and publish protocols, collaborate in communities, share insights through comments, and track progress with run records.

Create free account

OPEN  ACCESS



DOI: <https://dx.doi.org/10.17504/protocols.io.7zdhp26>

External link: <https://doi.org/10.1371/journal.pone.0225863>

Protocol Citation: Norfitriah Mohamed Sohaimi, Mohd Hair Bejo, Abdul Rahman Omar, Aini Ideris, Nurulfiza Mat Isa 2019. Propagation and attenuation of FAdV isolate in CEL cells. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.7zdhp26>

**Manuscript citation:**

Sohaimi NM, Bejo MH, Omar AR, Ideris A, Isa NM (2019) Molecular characterization of fowl adenovirus isolate of Malaysia attenuated in chicken embryo liver cells and its pathogenicity and immunogenicity in chickens. PLoS ONE 14(12): e0225863. doi: [10.1371/journal.pone.0225863](https://doi.org/10.1371/journal.pone.0225863)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: October 06, 2019

Last Modified: October 06, 2019

Protocol Integer ID: 28421

Keywords: cel cell, attenuation of fadv, cel, cell, fadv

Materials

MATERIALS

 Fetal Bovine Serum **Gibco - Thermo Fisher Scientific Catalog #10270106**

A confluent monolayer was washed twice with serum free medium and inoculated with 0.1mL of homogenate liver embryos and labeled as first passage (CEL1). Infected flasks were incubated at 37°C for 60 minutes for virus adsorption and added with maintenance medium containing 2% fetal bovine serum (FBS) under 37°C incubator. The cells were observed daily under inverted microscope for cytopathic effect (CPE) for 3 days post-inoculation (pi). Flasks with prominent CPE were harvested by 3 times repeated frozen and thawed prior centrifugation at 216 x g for 10 minutes. Virus supernatant was collected and stored at -20°C prior inoculation for subsequence passages. Fresh confluent monolayer was prepared for each passage and inoculated with 0.1mL of viral supernatant and continued until 35th consecutive passage. For non-infected flasks, monolayer remained uninoculated and was used as control cells.

