FEB 28, 2021

Protein A- Protein A Sandwich ELISA

Angel A Justiz-Vaillant¹¹University of the West Indies St. Augustine

Angel A Justiz-Vaillant

University of the West Indies St. Augustine

ABSTRACT

This ELISA was used to study Staphylococcal protein-A (SpA) interactions with various mammalian and avian immunoglobulins.

OPEN ACCESS**DOI:**

dx.doi.org/10.17504/protocols.io.bs4neyw

Protocol Citation: Angel A Justiz-Vaillant 2021. Protein A-Protein A Sandwich ELISA.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.bs4neyw>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Feb 28, 2021**Last Modified:** Feb 28, 2021**PROTOCOL integer ID:**

47740

- 1 This ELISA was used to study Staphylococcal protein-A (SpA) interactions with various mammalian and avian immunoglobulins. The 96 well microtiter plate was coated overnight at 4°C with 2 µg/µl per well of SpA in carbonate-bicarbonate buffer pH 9.6.

- 2 The plate was then treated with bovine serum albumin solution and washed 4X with PBS-Tween. Fifty microliters of immunoglobulins (1 mg/mL) was added and incubated for 1.30h at room temperature, and the microplate was then rewashed 4X with PBS-Tween.
- 3 Then, 50 μ L of peroxidase-labeled-SpA conjugate diluted 1:5000 in PBS-non-fat milk was added to each well and incubated for 1.30h at RT. The plate was washed 4X with PBS-Tween.
- 4 Fifty microliters of 3,3',5,5' - tetramethylbenzidine (TMB; Sigma-Aldrich) was pipetted into each well. The reaction was stopped with 50 μ L of a 3M H₂S O₄ solution.
- 5 The plate was visually assessed for color development and read on a microplate reader at 450 nm. A cut-off point was calculated as the mean of the optical density of the negative controls multiplied by two. The higher the OD value, the higher the affinity of SpA for avian immunoglobulins. The cut-off point was set to 0.30.