OCT 20, 2019

Protein expression in OnePot PURE cell-free system

Konstantinos Ragios¹¹EPFL - EPF Lausanne

iGEM EPFL

Konstantinos Ragios

EPFL - EPF Lausanne

ABSTRACT

In this protocol we explain the procedure of protein expression in the OnePot PURE cell-free system.

OPEN ACCESS

DOI:

dx.doi.org/10.17504/protocols.io.8avhse6

External link:

<https://pubs.acs.org/doi/10.1021/acssynbio.8b00427>

Protocol Citation: Konstantinos Ragios 2019. Protein expression in OnePot PURE cell-free system.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.8avhse6>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Oct 16, 2019

Last Modified: Oct 20, 2019

PROTOCOL integer ID:

28725

Keywords: iGEM, cell-free, OnePot, PURE, protein, expression

MATERIALS

Material/Consumables:

- OnePot Protein solution
- OnePot Ribosome solution
- OnePot Energy solution
- DNA template
- Nuclease free water

	Component	Volume (μL)	Final concentration of the reaction (nM)
	Protein Solution	0.65	-
	Ribosome Solution	0.9	-
	Energy Solution	2	-
	DNA Template	x	5
	Water	1.45-x	-

BEFORE START INSTRUCTIONS

In order to create the components needed refer to the following protocols:

Protocol

NAME

Protein Purification for OnePot PURE cell-free system

CREATED BY

Konstantinos Ragios

[PREVIEW](#)

Protocol

NAME

Ribosome Purification for OnePot PURE cell-free system

CREATED BY

Konstantinos Ragios

[PREVIEW](#)

Protocol

NAME

Energy solution preparation for OnePot PURE cell-free system

CREATED BY

Konstantinos Ragios

[PREVIEW](#)

Protein expression

- 1 For a 5µl reaction add to a tube 2µl of Energy solution, 0.65µl of Protein solution, 0.9µl of Ribosome solution, the DNA template (5nM final concentration in the reaction) and if needed add

water to reach the final volume.

Note

The minimum reaction volume is 5µl while the suggested one is 10µl.
For any final reaction volume (e.g. X µl) you just need to multiply the volumes of the components needed by a factor of X/5.

Note

If you haven't produce ribosomes then any store-bought product can be used.

2 Centrifuge for a few seconds and then your solution will start reacting.

3

Note

If you want to measure it with a plate reader, ideally you should add each component in different corners of the well, so the reaction will not initiate before adding the plate on the reader. The centrifuge for a few seconds.